


## SOLVED EXAMPLES

**Ex. 1** The weight of sodium bromate required to prepare 55.5 mL of 0.672 N solution for cell reaction,



(A) 1.56 g                      (B) 0.9386 g                      (C) 1.23 g                      (D) 1.32 g

**Sol.** Meq. of  $\text{NaBrO}_3 = 55.5 \times 0.672 = 37.296$

Let weight of  $\text{NaBrO}_3 = W$


$$\therefore \frac{W}{M_{\text{NaBrO}_3}} \times 6 \times 1000 = 37.296 \text{ (equivalent weight = } M/6 \text{) of n-factor = 6}$$

$$\therefore \frac{M}{151} \times 6 \times 1000 = 37.296$$

$$\therefore W = 0.9386 \text{ g}$$

Hence, (B) is the correct answer.

**Ex. 2**  $\text{NaIO}_3$  reacts with  $\text{NaHSO}_3$  according to equation



The weight of  $\text{NaHSO}_3$  required to react with 100 mL of solution containing 0.68 g of  $\text{NaIO}_3$  is

(A) 5.2 g                      (B) 0.2143 g                      (C) 2.3 g                      (D) none of the above

**Sol.** Meq. of  $\text{NaHSO}_3 = \text{Meq. of NaIO}_3 = N \times V = \frac{0.68}{198} \times 6 \times 1000$  ( $\text{I}^{5+} + 6\text{e}^- \rightarrow \text{I}^-$ )

$$\therefore \frac{W_{\text{NaHSO}_3}}{M_{\text{NaHSO}_3}} \times 2 \times 1000 = \frac{0.68}{198} \times 6 \times 1000$$


$$W_{\text{NaHSO}_3} = \frac{0.68 \times 6 \times 100 \times 104}{198 \times 1000} = 0.2143$$

Hence (B) is the correct answer.

**Ex. 3** If 0.5 moles of  $\text{BaCl}_2$  is mixed with 0.1 moles of  $\text{Na}_3\text{PO}_4$ , the maximum amount of  $\text{Ba}_3(\text{PO}_4)_2$  that can be formed is

(A) 0.7 mol                      (B) 0.5 mol                      (C) 0.2 mol                      (D) 0.05 mol

**Sol.** Let us first solve this problem by writing the complete balanced reaction.



We can see that the moles of  $\text{BaCl}_2$  used are  $\frac{3}{2}$  times the moles of  $\text{Na}_3\text{PO}_4$ . Therefore, to react with

0.1 mol of  $\text{Na}_3\text{PO}_4$ , the moles of  $\text{BaCl}_2$  required would be  $0.1 \times \frac{3}{2} = 0.15$ . Since  $\text{BaCl}_2$  is 0.5 mol, we can conclude that


$\text{Na}_3\text{PO}_4$  is the limiting reagent. Therefore, moles of  $\text{Ba}_3(\text{PO}_4)_2$  formed is  $0.1 \times \frac{3}{2} = 0.05$  mol.

Hence, (D) is the correct answer.

**Ex. 4** A 0.1097 g sample of  $\text{As}_2\text{O}_3$  required 36.10 mL of  $\text{KMnO}_4$  solution for its titration. The molarity of  $\text{KMnO}_4$  solution is.

(A) 0.02                      (B) 0.04                      (C) 0.0122                      (D) 0.3

**Sol.**



## CHEMISTRY FOR JEE MAIN & ADVANCED

Let, molarity of  $\text{KMnO}_4$  solution be  $M$

$\therefore$  Eq. of  $\text{As}_2\text{O}_3 = \text{Eq. of KMnO}_4$  solution

$$\frac{0.1097}{198} \times 4 = \frac{36.10 \times M \times 5}{1000} \quad \left( \text{Equivalent weight } \text{As}_2\text{O}_3 = \frac{198}{4} \right)$$

Molarity = 0.0122 M

Hence, (C) is the correct answer.

**Ex. 5** In basic medium,  $\text{CrO}_4^{2-}$  oxidize  $\text{S}_2\text{O}_3^{2-}$  to form  $\text{SO}_4^{2-}$  and itself changes to  $\text{Cr}(\text{OH})_4^-$ . How many mL of 0.154 M  $\text{CrO}_4^{2-}$  are required to react with 40 mL of 0.246 M  $\text{S}_2\text{O}_3^{2-}$ ?

(A) 200 mL                      (B) 156.4 mL                      (C) 170.4 mL                      (D) 190.4 mL

**Sol.**  $40 \times 0.246 \times 8 = V \times 0.154 \times 3$  (Meq. of  $\text{S}_2\text{O}_3^{2-} = \text{Meq. of CrO}_4^{2-}$ )

$$\therefore V = 170.4 \text{ mL}$$

Hence, (C) is the correct answer.

**Ex. 6** 10 mL of 0.4 M  $\text{Al}_2(\text{SO}_4)_3$  is mixed with 20 mL of 0.6 M  $\text{BaCl}_2$ . Concentration of  $\text{Al}^{3+}$  ion in the solution will be.

(A) 0.266 M                      (B) 10.3 M                      (C) 0.1 M                      (D) 0.25 M

**Sol.**  $\text{Al}_2(\text{SO}_4)_3 + \text{BaCl}_2 \longrightarrow \text{BaSO}_4 \downarrow + \text{AlCl}_3$

Initial Meq.                       $10 \times 0.4 \times 6$                        $20 \times 0.6 \times 2$                       0                      0

$$= 24 \qquad \qquad \qquad = 24$$

Final Meq.                      0                      0                      24                      24

$$[\text{Al}^{3+}] = \frac{24}{30 \times 3} = 0.266 \text{ M}$$

Hence (A) is the correct answer.

**Ex. 7** 0.52 g of a dibasic acid required 100 mL of 0.2 N NaOH for complete neutralization.

The equivalent weight of acid is

(A) 26                      (B) 52                      (C) 104                      (D) 156

**Sol.** Meq. of Acid = Meq. of NaOH

$$\frac{0.52}{E} \times 1000 = 100 \times 0.2$$


$$\therefore E = 26$$

Hence (A) is the correct answer.

**Ex. 8** 34 g hydrogen peroxide is present in 1120 mL of Solution : This solution is called

(A) 10 volume                      (B) 20 volume                      (C) 30 volume                      (D) 32 volume

**Sol.** Wt. of  $\text{H}_2\text{O}_2$  in 1 mL =  $\frac{34}{1120}$  g



34 g of  $\text{H}_2\text{O}_2$  gives 11200 mL of  $\text{O}_2$  at STP

$$\therefore \frac{34}{1120} \text{ g of } \text{H}_2\text{O}_2 \equiv \frac{11200}{34} \times \frac{34}{1120} = 10 \text{ mL of } \text{O}_2 \text{ at STP.}$$



Hence, (A) is the correct answer.

## REDOX REACTION AND EQUIVALENT CONCEPT

**Ex. 9** The number of moles of  $\text{KMnO}_4$  that will be required to react with 2 mol of ferrous oxalate is

- (A)  $\frac{6}{5}$                       (B)  $\frac{2}{5}$                       (C)  $\frac{4}{5}$                       (D) 1

**Sol.**  $\text{Mn}^{7+} + 5\text{e}^- \rightarrow \text{Mn}^{2+}] \times 3$



3 moles of  $\text{KMnO}_4 = 5$  moles of  $\text{FeC}_2\text{O}_4$

$$\therefore 2 \text{ mol of ferrous oxalate} \equiv \frac{6}{5} \text{ mole of } \text{KMnO}_4$$

Hence, (A) is the correct answer.

**Ex. 10** The minimum quantity of  $\text{H}_2\text{S}$  needed to precipitate 63.5 g of  $\text{Cu}^{2+}$  will be nearly.

- (A) 63.5 g                      (B) 31.75 g                      (C) 34 g                      (D) 2.0 g

**Sol.** Meq. of  $\text{H}_2\text{S} = \text{Meq. of } \text{Cu}^{2+}$

$$\therefore \frac{W_{\text{H}_2\text{S}}}{34/2} \times 1000 = \frac{63.5}{63.5/2} \times 1000$$

$$\therefore W_{\text{H}_2\text{S}} = 34 \text{ g}$$

Hence, (C) is the correct answer.

**Ex. 11** Which of the following is / are correct?

- (A) g mole wt. = mol. wt. in g = wt. of  $6.02 \times 10^{23}$  molecules  
 (B) mole = N molecule =  $6.02 \times 10^{23}$  molecules  
 (C) mole = g molecules  
 (D) none of the above

**Sol.** (A), (B) and (C)

**Ex. 12** 8 g of  $\text{O}_2$  has the same number of molecules as


- (A) 7 g of CO                      (B) 14 g of CO                      (C) 28 g of CO                      (D) 11 g of  $\text{CO}_2$

**Sol.** (A) and (D)

**Ex. 13** What volume of 0.1 M  $\text{KMnO}_4$  is needed to oxidize 100 mg of  $\text{FeC}_2\text{O}_4$  in acid solution ?

- (A) 4.1 mL                      (B) 8.2 mL                      (C) 10.2 mL                      (D) 4.6 mL

**Sol.** Meq. of  $\text{KMnO}_4 = \text{Meq. of } \text{FeC}_2\text{O}_4$



$$\therefore V = 4.1 \text{ mL}$$

Hence, (A) is the correct answer.

## CHEMISTRY FOR JEE MAIN & ADVANCED


**Ex. 14** What volume of 6 M  $\text{HNO}_3$  is needed to oxidize 8 g of  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$ ,  $\text{HNO}_3$  gets converted to  $\text{NO}$ ?

- (A) 8 mL                      (B) 7.936 mL                      (C) 32 mL                      (D) 64 mL

**Sol.** Meq. of  $\text{HNO}_3 = \text{Meq. of Fe}^{2+}$

$$\text{or } 6 \times 3 \times V = \frac{8}{56} \times 1000$$

$$V = 7.936 \text{ mL}$$



Hence, (B) is the correct answer.

**Ex. 15** 0.5 g of fuming  $\text{H}_2\text{SO}_4$  (oleum) is diluted with water. This solution is completely neutralized by 26.7 mL of 0.4 N  $\text{KOH}$ . The percentage of free  $\text{SO}_3$  in the sample is

- (A) 30.6%                      (B) 40.6%                      (C) 20.6%                      (D) 50%

**Sol.** Meq. of  $\text{H}_2\text{SO}_4 + \text{Meq. of SO}_3 = \text{Meq. of KOH}$

$$\therefore \frac{(0.5 - x)}{98/2} \times 1000 + \frac{x}{80/2} \times 1000 = 26.7 \times 0.4$$

$$\therefore x = 0.103$$

$$\therefore \% \text{ of SO}_3 = \frac{0.103}{0.5} \times 100 = 20.6\%$$

Hence, (C) is the correct answer.

**Ex. 16** The eq. wt. of a substance is the weight which either combines or displaces.

- (A) 8 part of O                      (B) 2 part of H                      (C) 35.5 part of Cl                      (D) none of the above

**Sol.** (A) and (C)

**Ex. 17** 'A' g of a metal displaces V mL of  $\text{H}_2$  at NTP. Eq. wt of metal E is / are :

$$(A) E = \frac{A}{\text{Wt. of H}_2 \text{ displaced}} \times \text{Eq. wt. of H}$$




$$(B) E = \frac{A \times 1.008 \times 22400}{\text{Volume of H}_2 \text{ displaced} \times 2} \times \text{Eq. wt. of H}$$

$$(C) E = \frac{A \times 1.008}{\text{Volume of H}_2 \text{ displaced} \times 0.0000897}$$

(D) none of the above

**Sol.** (A), (B) and (C)

**Ex. 18** Which of the following is/are redox reaction (s)?



**Sol.** (B), (C) and (D)

**Ex. 19** Which of the following statements is/are true if 1 mol of  $\text{H}_3\text{PO}_x$  is completely neutralized by 40 g of  $\text{NaOH}$ ?

(A)  $x = 2$  and acid is monobasic


(B)  $x = 3$  and acid is dibasic

(C)  $x = 4$  and acid is tribasic

(D)  $x = 2$  and acid does not form acid salt

**Sol.** (A) and (D).

**Ex. 20** In the reaction,



(A) bromine is oxidized and carbonate is reduced

(B) bromine is oxidized

(C) bromine is reduced

(D) it is disproportionation reaction or autoredox change

**Sol.** (B), (C) and (D)

**Ex. 21** A 20 g sample of only CuS and Cu<sub>2</sub>S was treated with 100 mL of 1.25 M K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>. The products obtained were Cr<sup>3+</sup>, Cu<sup>2+</sup> and SO<sub>2</sub>. The excess oxidant was reacted with 50 mL of Fe<sup>2+</sup> Solution : 25 mL of the same Fe<sup>2+</sup> solution required 0.875 M KMnO<sub>4</sub> under acidic condition, the volume of KMnO<sub>4</sub> used was 20 mL. Find the % of CuS and Cu<sub>2</sub>S in the sample.

**Sol.** Equivalents of dichromate initially =  $\frac{1.25 \times 6 \times 100}{1000} = 0.75$

$$\text{Equivalents of Fe}^{2+} \text{ in 25 mL} = \frac{0.875 \times 5 \times 5 \times 20}{1000} = 0.0875$$

$$\text{Equivalents of Fe}^{2+} \text{ in 50 mL} = 0.0875 \times 2 = 0.175$$

$$\text{Equivalents of excess dichromate} = 0.175$$

$$\therefore \text{Equivalents of dichromate consumed by (CuS and Cu}_2\text{S)} \\ = 0.75 - 0.175 = 0.575$$

If x g is the mass of CuS, the mass of Cu<sub>2</sub>S is (10 - x) g

$$\frac{x}{95.5} \times 6 + \frac{(10-x)}{159} \times 8 = 0.575$$

$$\therefore x = 5.74 \text{ g}$$


$$\% \text{ CuS} = \frac{5.74}{20} \times 100 = 28.7 \%$$

$$\% \text{ Cu}_2\text{S} = 71.3 \%$$



**Ex. 22** 2.249 g of a sample of pure BaCO<sub>3</sub> and impure CaCO<sub>3</sub> containing some CaO was treated with dil. HCl and it evolved 168 mL of CO<sub>2</sub> at N.T.P. From this solution, BaCrO<sub>4</sub> was precipitated filtered and washed. The precipitate was dissolved in dil. H<sub>2</sub>SO<sub>4</sub> and diluted to 100 mL. 10 mL of this solution when treated with KI solution liberated iodine which required exactly 20 mL of 0.05 N Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>. Calculate the percentage of CaO in the sample.

**Sol.** Let, weight of BaCO<sub>3</sub>, CaCO<sub>3</sub> and CaO are x, y and z respectively.

$$\therefore x + y + z = 2.249$$



Redox change



$$\text{Meq. of BaCO}_3 = \text{Meq. of BaCrO}_4 = \text{Meq. of I}_2$$

$$\therefore \frac{x}{197/3} \times 1000 = 20 \times 0.05 \times \frac{100}{10}$$

$$\therefore x = 0.657 \text{ g}$$

...(1)

The equivalent weight of BaCrO<sub>4</sub> is M/3, therefore for BaCO<sub>3</sub>, it should be M/3 also because mole ratio of BaCO<sub>3</sub> and BaCrO<sub>4</sub> is 1 : 1.

Applying POAC for C atom,

$$\text{Moles of C in BaCO}_3 + \text{Moles of C in CaCO}_3 = \text{Moles of C in CO}_2$$

$$\begin{aligned} \therefore \quad \frac{x}{197} + \frac{y}{100} &= \frac{168}{22400} \\ \Rightarrow \quad 200x + 294y &= 295.5 \quad \dots(2) \\ \text{From equation (1) and (2)} \\ y &= 0.416 \text{ g} \\ \therefore \quad 0.657 + 0.416 + z &= 2.249 \\ z &= 1.176 \\ \% \text{ of CaO} &= \frac{1.176}{2.249} = 52.29\% \end{aligned}$$

**Ex. 23** For estimating ozone in the air, a certain volume of air is passed through an alkaline KI solution when  $O_2$  is evolved and iodide is oxidized to iodine. When such a solution is acidified, free iodine is evolved which can be titrated with standard  $Na_2S_2O_3$  solution : In an experiment, 10 L of air at 1 atm and  $27^\circ C$  were passed through an alkaline KI solution, and at the end, the iodine was entrapped in a solution which on titration as above required 1.5 mL of 0.02 N  $Na_2S_2O_3$  solution. Calculate volume percentage of ozone in the sample.

**Sol.** The chemical reaction is,  $H_2O + KI + O_3 \longrightarrow I_2 + O_2 + KOH$   
 Milliequivalents of iodine = Milliequivalents of KI = Milliequivalents of  $O_3$  reacted  
 Milliequivalents of  $Na_2S_2O_3 = 1.5 \times 0.02 = 3 \times 10^{-2}$   
 Millimoles of iodine =  $\frac{3 \times 10^{-2}}{2} = 1.5 \times 10^{-2}$  [ $\because$  n-factor for iodine = 2]  
 Millimoles of ozone =  $1.5 \times 10^{-2}$   
 Volume of ozone =  $\frac{nRT}{P} = \frac{1.5 \times 10^{-5} \times 0.0821 \times 300}{1} = 36.945 \times 10^{-5}$  litre  
 Volume percent of ozone =  $\frac{36.945 \times 10^{-5}}{10} \times 100 = 3.6945 \times 10^{-3}$

**Ex. 24** What volume of 0.40 M  $H_2SO_4$  is required to produce 34.0 g of  $H_2S$  by the reaction,  
 $8KI + 5H_2SO_4 \rightarrow 4K_2SO_4 + 4I_2 + H_2S + 4H_2O$ ?

**Sol.** 1 mole of  $H_2S \equiv 5$  moles of  $H_2SO_4$   
 34 g of  $H_2S \equiv 5$  moles of  $H_2SO_4$   
 $0.40 \times V_{H_2SO_4} = 5$   
 $\therefore V_{H_2SO_4} = \frac{5}{0.40} = 12.5$  litre

**Ex. 25** To 50 L of 0.2 N NaOH, 2.5 L of 2N HCl and 15 L of 0.1 N  $FeCl_3$  solutions are added. What weight of  $Fe_2O_3$  can be obtained from the precipitate? Also report the normality of NaOH left in resultant solution :

**Sol.** Eq. of NaOH =  $50 \times 0.2 = 10$   
 Eq. of HCl =  $2.5 \times 2 = 5$   
 Eq. of NaOH left after reaction with HCl =  $10 - 5 = 5$   
 $FeCl_3 + NaOH \rightarrow Fe(OH)_3 \downarrow \xrightarrow{\Delta} Fe_2O_3$   
 $FeCl_3$  reacts with NaOH to give  $Fe(OH)_3$  which on ignition gives  $Fe_2O_3$   
 $\therefore$  Eq. of NaOH used for  $FeCl_3 =$  Eq. of  $Fe(OH)_3$   
 $=$  Eq. of  $Fe_2O_3$   
 $= 15 \times 0.1 = 3.5$

$$\therefore \text{Eq. of NaOH left finally} = 5 - 1.5 = 3.5$$

$$\text{Normality of NaOH in the resultant solution} = \frac{3.5}{70} = 0.05$$

$$\frac{W_{\text{Fe}_2\text{O}_3}}{M_{\text{Fe}_2\text{O}_3}} \times 6 = 1.5 \quad (\text{n-factor for Fe}_2\text{O}_3 = 6)$$

$$W_{\text{Fe}_2\text{O}_3} = \frac{1.5 \times 160}{6} = 40\text{g}$$

**Ex. 26** One litre of a mixture of  $\text{O}_2$  and  $\text{O}_3$  at NTP was allowed to react with an excess of acidified solution of KI. The iodine liberated required 30 mL of M/10  $\text{Na}_2\text{S}_2\text{O}_3$  solution for titration. What is the weight percent of ozone in the mixture? Ultraviolet radiation of wavelength 300 nm can decompose ozone. Assuming that one photon can decompose one ozone molecule, how many photons would have been required for the complete decomposition of ozone in the original mixture?

**Sol.**  $\text{O}_3 + 2\text{KI} + \text{H}_2\text{O} \rightarrow 2\text{KOH} + \text{I}_2 + \text{O}_2$   
 $2\text{Na}_2\text{S}_2\text{O}_3 + \text{I}_2 \rightarrow \text{Na}_2\text{S}_4\text{O}_6 + 2\text{NaI}$   
 Millimoles of  $\text{O}_3 = \text{Millimoles of I}_2 = \frac{1}{2} \times \text{Millimoles of Na}_2\text{S}_2\text{O}_3$   
 $= \frac{1}{2} \times 30 \times 1/10 = 1.5 \text{ millimoles}$   
 $= 0.0015 \text{ moles}$

Total moles of  $\text{O}_3$  and  $\text{O}_2$  in the mixture

$$PV = nRT$$

$$1 \times 1 = n \times 0.0821 \times 273$$

$$\therefore n = 0.044$$

$$\text{Moles of O}_2 = 0.044 - 0.0015 = 0.0425$$

$$W_{\text{O}_2} = 0.0425 \times 32 \text{ g} = 1.36 \text{ g}$$

$$W_{\text{O}_3} = 0.0015 \times 48 \text{ g} = 0.072 \text{ g}$$

$$\therefore \% \text{ of O}_3 = \frac{0.072}{1.432} \times 100 = 5.028 \%$$

$$\text{No. of photons required for decomposition of Ozone molecules} = 0.0015 \times 6.023 \times 10^{23} = 0.90 \times 10^{21}$$

**Ex. 27** A 2.18 g sample contains a mixture of XO and  $\text{X}_2\text{O}_3$ . It reacts with 0.015 moles of  $\text{K}_2\text{Cr}_2\text{O}_7$  to oxidize the sample completely to form  $\text{XO}_4^-$  and  $\text{Cr}^{3+}$ . If 0.0187 mole of  $\text{XO}_4^-$  is formed, what is the atomic mass of X?

**Sol.**  $\text{XO} + \text{K}_2\text{Cr}_2\text{O}_7 \longrightarrow \text{Cr}^{3+} + \text{XO}_4^-$   
 $\text{X}_2\text{O}_3 + \text{K}_2\text{Cr}_2\text{O}_7 \longrightarrow \text{Cr}^{3+} + \text{XO}_4^-$   
 Let, wt. of XO in the mixture be x g  
 Equivalent of  $\text{K}_2\text{Cr}_2\text{O}_7$  consumed by the mixture =  $0.015 \times 6$

$$\text{Equivalent of XO} = \frac{x}{x+16} \times 5$$

$$\text{Equivalent of X}_2\text{O}_3 = \frac{2.18 - x}{2x + 48} \times 8$$

$$\therefore \frac{x}{x+16} \times 5 + \frac{2.18 - x}{2x - 148} \times 8 = 0.015 \times 6$$

## CHEMISTRY FOR JEE MAIN & ADVANCED

Since 1 mole of XO gives 1 mole  $\text{XO}_4^-$  and 1 mole of  $\text{X}_2\text{O}_3$  gives 2 moles of  $\text{XO}_4^-$

$$\therefore \frac{x}{x+16} + \frac{2x(2.18-x)}{2x+48} = 0.0187$$

On solving,  $x = 99$

**Ex. 28** An aqueous solution containing 0.10 g  $\text{KIO}_3$  (formula wt. = 214.0) was treated with an excess of KI Solution : The solution was acidified with HCl. The liberated  $\text{I}_2$  consumed 55 mL of thiosulphate solution to decolourize the blue starch-iodine complex. Calculate molarity of the sodium thiosulphate solution :

**Sol.** Moles of  $\text{KIO}_3 = \frac{0.1}{214} = 0.00047$

$$\therefore \text{Moles of } \text{I}_2 \text{ liberated from } \text{KIO}_3 = \frac{0.00047}{2} = 0.000235$$


$$\text{Moles of KI reacting} = 0.00047 \times 5 = 0.00235 \quad (\because \text{n-factor for } \text{KIO}_3 \text{ and KI are 5 and 1 respectively})$$

$$\text{Moles of } \text{I}_2 \text{ produced from KI} = \frac{0.00235}{2} = 0.001175$$

Total moles of  $\text{I}_2$  produced and reacted =  $0.000235 + 0.001175 = 0.00141$  equivalents of  $\text{I}_2$  reacted =  $0.00141 \times 2 = 0.00282$  = equivalents of thiosulphate Solution :

$$\text{Molarity} = \frac{0.00282 \times 1000}{55} = 0.0512 \text{ M (For thiosulphate n-factor} = 1)$$

**Ex. 29** Mg can reduce  $\text{NO}_3^-$  to  $\text{NH}_3$  in basic medium.




A 25.0 mL sample of  $\text{NO}_3^-$  solution was treated with Mg. The  $\text{NH}_3$  (g) was passed into 100 mL of 0.15 N HCl. The excess of HCl required 32.10 mL of 0.10 N NaOH for neutralization. What was the molarity of  $\text{NO}_3^-$  ions in the original sample ?

**Sol.** Meq. of  $\text{NH}_3$  formed = Meq. of HCl used  
 $= 100 \times 0.15 - 32.10 \times 0.10$   
 $= 11.79$

Here, n-factor of  $\text{NH}_3$  is 1 (acid - base reaction)

For redox change,





$$\therefore \text{Meq. of } \text{NH}_3 \text{ for n-factor } 8 = 8 \times 11.79$$

$$\therefore \text{Normality of } \text{NO}_3^- = \frac{94.32}{25} = 3.77$$

$$\text{Molarity of } \text{NO}_3^- = \frac{3.77}{8} = 0.47125$$



**Ex. 30** Chile salt peter a source of  $\text{NaNO}_3$  also contains  $\text{NaIO}_3$ . The  $\text{NaIO}_3$  can be used as a source of iodine produced in the following reactions :



One litre of chile salt peter solution containing 6.80 g  $\text{NaIO}_3$ , is treated with stoichiometric quantity of  $\text{NaHSO}_3$ . Now additional amount of same solution is added to the reaction mixture to bring about the second reaction. How many grams of  $\text{NaHSO}_3$  are required in step 1 and what additional volume of chile salt peter must be added in step 2 to bring in complete conversion of  $\text{I}^-$  to  $\text{I}_2$  ?

**Sol.** Meq. of  $\text{NaHSO}_3 = \text{Meq. of NaIO}_3 = N \times V = \frac{6.8}{198} \times 6 \times 1000$



$$\therefore \text{Meq. of NaHSO}_3 = 206.06$$

$$\frac{W_{\text{NaHSO}_3}}{M/2} \times 1000 = 206.06$$

$$\Rightarrow W_{\text{NaHSO}_3} = \frac{206.06 \times 104}{2 \times 1000} = 10.715 \text{ g}$$

Also, Meq. of  $\text{I}^-$  formed using n-factor 6 = 206.06

In step 2, n-factor  $\text{I}^-$  is 1 and n-factor for  $\text{IO}_3^-$  is 5


$$\text{Meq. of I}^- \text{ formed using n-factor 1} = \frac{206.06}{6}$$

$$\text{Meq. of NaIO}_3 \text{ used in 2}^{\text{nd}} \text{ step} = \frac{206.06}{6}$$

$$\Rightarrow \frac{6.8}{198} \times 5 \times V = \frac{206.06}{6}$$

$$\therefore V_{\text{NaIO}_3} = 199.99 \text{ mL}$$



**Ex. 31** 1 g sample of  $\text{AgNO}_3$  is dissolved in 50 mL of water. It is titrated with 50 mL of KI Solution : The  $\text{AgI}$  precipitated is filtered off. Excess of KI in filtrate is titrated with M/10  $\text{KIO}_3$  in presence of 6 M  $\text{HCl}$  till all  $\text{I}^-$  converted into  $\text{ICl}$ . It requires 50 ml of M/10  $\text{KIO}_3$  Solution : 20 mL of the same stock solution of KI require 30 mL of M/10  $\text{KIO}_3$  under similar conditions. Calculate % of  $\text{AgNO}_3$  in the sample. The reaction is :



**Sol.**  $\text{AgNO}_3 + \text{KI} \longrightarrow \text{AgI} \downarrow + \text{KNO}_3$

As present in  $\text{AgNO}_3$  is removed as  $\text{AgI}$  by adding 50 mL of KI solution of which 20 mL requires 30 mL of M/10  $\text{KIO}_3$

$$\text{Meq. of KI in 20 mL} = \text{Meq. of KIO}_3 = 30 \times \frac{1}{10} \times 4$$



$$\therefore \text{Eq. wt. of KI} = \frac{M}{2}$$

$$\therefore \text{Meq. of KI in 50 mL added to AgNO}_3 = \frac{30 \times 4 \times 50}{10 \times 20} = 30$$

$$\text{Meq. of KI left unused by AgNO}_3 = 50 \times \frac{1}{10} \times 4 = 20$$

$$\therefore \text{Meq. of KI used for AgNO}_3 = 30 - 20 = 10$$

$$\therefore \text{Meq. of AgNO}_3 = 10$$

$$\frac{W_{\text{AgNO}_3}}{170} \times 2 \times 1000 = 10$$

$$\therefore W_{\text{AgNO}_3} = 0.85 \text{ g} = 85\%$$


$$\frac{\text{Mole of AgNO}_3}{\text{Mole of KI}} = \frac{1}{1}$$

$$\therefore \text{If equivalent weight of KI} = \frac{M}{2}$$

$$\text{Equivalent weight of AgNO}_3 = \frac{M}{2}$$


**Ex. 32** A sample of Mg was burnt in air to give a mixture of MgO and Mg<sub>3</sub>N<sub>2</sub>. The ash was dissolved in 60 Meq. of HCl and the resulting solution was back titrated with NaOH. 12 Meq. of NaOH were required to reach the end point. An excess of NaOH was then added and the solution was distilled. The ammonia released was then trapped in 10 Meq. of acid. Back titration of this solution required 6 Meq. of base. Calculate the % of Mg burnt to the nitride.

**Sol.** Let, total moles of Mg used for MgO and Mg<sub>3</sub>N<sub>2</sub> be a and b respectively



$$a \qquad \qquad \qquad 0$$


$$0 \qquad \qquad \qquad a$$



$$b \qquad \qquad \qquad 0$$

$$0 \qquad \qquad \qquad b/3$$

Now a,  $\frac{b}{3}$  moles respectively of MgO and Mg<sub>3</sub>N<sub>2</sub> are present in the mixture.



or solution contains (A) moles of MgCl<sub>2</sub> from MgO and (B) moles of MgCl<sub>2</sub> from Mg<sub>3</sub>N<sub>2</sub> and  $\frac{2b}{3}$  moles of NH<sub>4</sub>Cl.

Now, moles of HCl or Meq. of HCl (monobasic) = 30 - 12 = 48

$$2a + \frac{8b}{3} = 48 \qquad \qquad \qquad \dots\dots\dots(1)$$

Moles of NH<sub>4</sub>Cl formed = Moles of NH<sub>3</sub> liberated = Moles of HCl used for absorbing NH<sub>3</sub>.

$$\frac{2b}{3} = (10 - 6) = 4 \qquad \qquad \qquad \dots\dots\dots(2)$$

$$\text{From (1), } 2a + \frac{8 \times 6}{3} = 48 \text{ or } a = 16$$

$$\therefore \% \text{ of Mg used for Mg}_3\text{N}_2 = \frac{6}{(6+16)} \times 100 = 27.27\%$$


## REDOX REACTION AND EQUIVALENT CONCEPT

**Ex. 33** 30 mL of a solution containing 9.15 g/litre of an oxalate  $K_xH_y(C_2O_4)_z \cdot nH_2O$  are required for titrating 27 mL of 0.12 N NaOH and 36 mL of 0.12 N  $KMnO_4$  separately. Calculate x, y, z and n. Assume all H atoms (except  $H_2O$ ) are replaceable and x, y, z are in the simple ratio of g atoms.

**Sol.** Let molecular weight of oxalate salt is M

(i) n - factor in acid - base reaction = 2

(ii) n - factor in redox titration =  $2 \times z$



$\therefore$  Meq. of acid in 30 mL = Meq. of NaOH used

$$30 \times \frac{9.15}{M} \times y = 27 \times 0.12 \quad \dots(1)$$

$$\text{Also, } 30 \times \frac{9.15}{M} \times (2z) = 36 \times 0.12 \quad \dots(2)$$

$$\text{From equations (1) and (2) } \frac{y}{2z} = \frac{27}{36} \Rightarrow \frac{y}{z} = \frac{3}{2} \quad \dots(3)$$

Also, total cationic charge = total anionic charge

$$\therefore x + y = 2z \quad \dots(4)$$

By equations (3) and (4)

$$x : y : z :: 1 : 3 : 2$$

These are in simplest ratio and molecular formula is  $KH_3(C_2O_4)_2 \cdot nH_2O$

Molecular weight of salt =  $39 + 3 + 176 + 18n = 218 + 18n$

$$\text{From equation (1), } M = \frac{30 \times 9.15 \times 3}{27 \times 0.12} = 254.16$$

$$\therefore 218 + 18n = 254.15$$


$$\therefore n = 2$$

$\therefore$  Oxalate salt is  $KH_3(C_2O_4)_2 \cdot 2H_2O$


**Ex. 34** Hydrogen peroxide solution (20 mL) reacts quantitatively with a solution of  $KMnO_4$  (20 mL) acidified with dilute  $H_2SO_4$ . The same volume of the  $KMnO_4$  solution is just decolourized by 10 mL of  $MnSO_4$  in neutral medium simultaneously forming a dark brown precipitate of hydrated  $MnO_2$ . The brown precipitate is dissolved in 20 mL of 0.2 M sodium oxalate under boiling condition in the presence of dilute  $H_2SO_4$ . Write the balanced equations involved in the reactions and calculate the volume strength of  $H_2O_2$ .

**Sol.** Meq. of  $MnO_2$  = Meq. of  $Na_2C_2O_4$

$$= 20 \times 0.2 \times 2 = 8$$



$$\text{Millimoles of } MnO_2 = \frac{8}{2} = 4$$






(Mole ratio is reciprocal of n-factor ratio)

$$\frac{\text{Millimoles of MnO}_4^-}{\text{Millimoles of MnO}_2} = \frac{2}{5}$$

$$\begin{aligned} \text{Millimoles MnO}_4^- &= \frac{2}{5} \times \text{Millimoles of MnO}_2 \\ &= \frac{2}{5} \times 4 = \frac{8}{5} \end{aligned}$$



$$\frac{\text{Millimoles of H}_2\text{O}_2}{\text{Millimoles of MnO}_4^-} = \frac{5}{2}$$

$$\therefore \text{Millimoles of H}_2\text{O}_2 = \frac{5}{2} \times \frac{8}{5} = 4$$



$$N_{\text{H}_2\text{O}_2} \times 20 = 4 \times 2 \quad (\text{n-factor for H}_2\text{O}_2 = 2)$$

$$\therefore N_{\text{H}_2\text{O}_2} = 0.4$$

$$\begin{aligned} \text{Volume strength of H}_2\text{O}_2 &= 5.6 \times N_{\text{H}_2\text{O}_2} \\ &= 5.6 \times 0.4 = 2.24 \end{aligned}$$

**Ex. 35** 1.0 g of moist sample of mixture of potassium chlorate ( $\text{KClO}_3$ ) and potassium chloride ( $\text{KCl}$ ) was dissolved in water and solution made upto 250 mL. This solution was treated with  $\text{SO}_2$  to reduce all  $\text{ClO}_3^-$  to  $\text{Cl}^-$  and excess of  $\text{SO}_2$  was removed by boiling. The total chloride was precipitated as silver chloride. The weight of precipitate was found to be 1.435 g. In another experiment, 25 mL of the original solution was heated with 30 ml 0.2 N  $\text{FeSO}_4$  and unused  $\text{FeSO}_4$  required 37.5 mL of 0.08 N  $\text{KMnO}_4$  solutions.

Calculate the molar ratio of the  $\text{ClO}_3^-$  to the given mixture.



**Sol.**  $\text{ClO}_3^-$  is reduced to  $\text{Cl}^-$  by  $\text{SO}_2$  and  $\text{ClO}_3^-$  is also reduced to  $\text{Cl}^-$  by  $\text{Fe}^{2+}$ , hence  $\text{AgCl}$  is formed due to total  $\text{Cl}^-$

$$\text{Meq. of Fe}^{2+} \text{ initially taken} = 30 \times 0.2 = 6$$

$$\text{Meq. of Fe}^{2+} \text{ unused} = 37.5 \times 0.08 = 3$$

$$\therefore \text{Meq. of Fe}^{2+} = 6.0 - 3.0 = 3.0$$

$$\text{Thus, Meq. of ClO}_3^- \text{ in 25 mL} = 3.0$$

$$\text{Moles of ClO}_3^- \text{ in 25 mL} = \frac{3.0}{1000 \times 6} = 0.0005$$



$$\text{O.N.} \quad \begin{array}{ccc} & 5 & -1 \end{array}$$

$$\text{Thus, moles of ClO}_3^- \text{ in 25 mL solution} = 0.0005$$

$\text{ClO}_3^-$  is also reduced to  $\text{Cl}^-$  by  $\text{SO}_2$  in first experiment and precipitated as  $\text{AgCl}$ .

$$\text{Thus, Cl}^- \text{ formed from ClO}_3^- = \text{AgCl from ClO}_3^- = 1.435 \text{ g}$$

$$\text{Total AgCl formed both from actual and Cl}^- \text{ from ClO}_3^- = 1.435 \text{ g}$$

$$= \frac{1.435}{143.5} = 0.01 \text{ mol}$$

$$\text{Thus, AgCl formed due to Cl}^- \text{ only} = 0.01 - 0.0005 = 0.0095 \text{ mol}$$

$$\text{Thus, ClO}_3^- \text{ and Cl}^- \text{ are in molar ratio} = \mathbf{1 : 19}$$

## Exercise # 1

[Single Correct Choice Type Questions]

1. Which reaction does not represent autoredox or disproportionation :-
- (A)  $\text{Cl}_2 + \text{OH}^- \longrightarrow \text{Cl}^- + \text{ClO}_3^- + \text{H}_2\text{O}$       (B)  $2\text{H}_2\text{O}_2 \longrightarrow \text{H}_2\text{O} + \text{O}_2$   
 (C)  $2\text{Cu}^+ \longrightarrow \text{Cu}^{2+} + \text{Cu}$       (D)  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7 \longrightarrow \text{N}_2 + \text{Cr}_2\text{O}_3 + 4\text{H}_2\text{O}$
2. Match List-I (Compounds) with List-II (Oxidation states of nitrogen) and select answer using the codes given below the lists :-
- |     | List-I                 |    | List-II |
|-----|------------------------|----|---------|
| (A) | $\text{NaN}_3$         | 1. | +5      |
| (B) | $\text{N}_2\text{H}_2$ | 2. | +2      |
| (C) | $\text{NO}$            | 3. | -1/3    |
| (D) | $\text{N}_2\text{O}_5$ | 4. | -1      |
- Code : (A) (B) (C) (D)
- |     |   |   |   |   |
|-----|---|---|---|---|
| (A) | 3 | 4 | 2 | 1 |
| (B) | 4 | 3 | 2 | 1 |
| (C) | 3 | 4 | 1 | 2 |
| (D) | 4 | 3 | 1 | 2 |
3. Which of the following is a redox reaction :-
- (A)  $2\text{CrO}_4^{2-} + 2\text{H}^+ \rightarrow \text{Cr}_2\text{O}_7^{2-} + \text{H}_2\text{O}$       (B)  $\text{CuSO}_4 + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4]\text{SO}_4$   
 (C)  $\text{Na}_2\text{S}_2\text{O}_3 + \text{I}_2 \rightarrow \text{Na}_2\text{S}_4\text{O}_6 + \text{NaI}$       (D)  $\text{Cr}_2\text{O}_7^{2-} + 2\text{OH}^- \rightarrow 2\text{CrO}_4^{2-} + \text{H}_2\text{O}$
4. 50 mL of 0.1 M solution of a salt reacted with 25 mL of 0.1 M solution of sodium sulphite. The half reaction for the oxidation of sulphite ion is :-
- $$\text{SO}_3^{2-}(\text{aq}) + \text{H}_2\text{O}(\ell) \longrightarrow \text{SO}_4^{2-}(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^-$$
- If the oxidation number of metal in the salt was 3, what would be the new oxidation number of metal :
- (A) zero      (B) 1      (C) 2      (D) 4
5. An element A in a compound ABD has oxidation number  $A^{n-}$ . It is oxidised by  $\text{Cr}_2\text{O}_7^{2-}$  in acid medium. In the experiment  $1.68 \times 10^{-3}$  moles of  $\text{K}_2\text{Cr}_2\text{O}_7$  were used for  $3.26 \times 10^{-3}$  moles of ABD. The new oxidation number of A after oxidation is
- (A) 3      (B)  $3 - n$       (C)  $n - 3$       (D)  $+n$
6. The incorrect order of decreasing oxidation number of S in compounds is :-
- (A)  $\text{H}_2\text{S}_2\text{O}_7 > \text{Na}_2\text{S}_4\text{O}_6 > \text{Na}_2\text{S}_2\text{O}_3 > \text{S}_8$       (B)  $\text{H}_2\text{SO}_5 > \text{H}_2\text{SO}_3 > \text{SO}_2 > \text{H}_2\text{S}$   
 (C)  $\text{SO}_3 > \text{SO}_2 > \text{H}_2\text{S} > \text{S}_8$       (D)  $\text{H}_2\text{SO}_4 > \text{SO}_2 > \text{H}_2\text{S} > \text{H}_2\text{S}_2\text{O}_8$
7. In which of the following reaction is there a change in the oxidation number of nitrogen atoms :-
- (A)  $2\text{NO}_2 \rightarrow \text{N}_2\text{O}_4$       (B)  $\text{NH}_3 + \text{H}_2\text{O} \rightarrow \text{NH}_4^+ + \text{OH}^-$   
 (C)  $\text{N}_2\text{O}_5 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3$       (D) none
8. In the reaction
- $$x\text{HI} + y\text{HNO}_3 \longrightarrow \text{NO} + \text{I}_2 + \text{H}_2\text{O}$$
- (A)  $x = 3, y = 2$       (B)  $x = 2, y = 3$       (C)  $x = 6, y = 2$       (D)  $x = 6, y = 1$
9. For the redox reaction :
- $$\text{MnO}_4^- + \text{C}_2\text{O}_4^{2-} + \text{H}^+ \longrightarrow \text{Mn}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$$
- the correct stoichiometric coefficients of  $\text{MnO}_4^-$ ,  $\text{C}_2\text{O}_4^{2-}$  and  $\text{H}^+$  are respectively
- (A) 2,5,16      (B) 16,5,2      (C) 5,16,2      (D) 2,16,5

## CHEMISTRY FOR JEE MAIN & ADVANCED

10. A certain weight of pure  $\text{CaCO}_3$  is made to react completely with 200 mL of an HCl solution to give 224 mL of  $\text{CO}_2$  gas at STP. The normality of the HCl is :-  
 (A) 0.05 N (B) 0.1 N (C) 1.0 N (D) 0.2 N
11. The volume of 1.5 M  $\text{H}_3\text{PO}_4$  solution required to neutralize exactly 90 mL of a 0.5 M  $\text{Ba}(\text{OH})_2$  solution is :-  
 (A) 10 mL (B) 30 mL (C) 20 mL (D) 60 mL
12. Volume  $V_1$  mL of 0.1 M  $\text{K}_2\text{Cr}_2\text{O}_7$  is needed for complete oxidation of 0.678 g  $\text{N}_2\text{H}_4$  in acidic medium. The volume of 0.3 M  $\text{KMnO}_4$  needed for same oxidation in acidic medium will be :-  
 (A)  $\frac{2}{5}V_1$  (B)  $\frac{5}{2}V_1$  (C)  $113V_1$  (D) can't say
13. Which of the following relations is incorrect :-  
 (A)  $3\text{N Al}_2(\text{SO}_4)_3 = 0.5\text{M Al}_2(\text{SO}_4)_3$  (B)  $3\text{M H}_2\text{SO}_4 = 6\text{N H}_2\text{SO}_4$   
 (C)  $1\text{M H}_3\text{PO}_4 = 1/3\text{N H}_3\text{PO}_4$  (D)  $1\text{M Al}_2(\text{SO}_4)_3 = 6\text{N Al}_2(\text{SO}_4)_3$
14. The mass of oxalic acid crystals ( $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ ) required to prepare 50 mL of a 0.2 N solution is :-  
 (A) 4.5 g (B) 6.3 g (C) 0.63 g (D) 0.45 g
15. 125 mL of 63% (w/v)  $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$  is made to react with 125 mL of a 40% (w/v) NaOH solution. The resulting solution is :-  
 (A) neutral (B) acidic (C) strongly acidic (D) alkaline
16. If equal volumes of 0.1 M  $\text{KMnO}_4$  and 0.1 M  $\text{K}_2\text{Cr}_2\text{O}_7$  solutions are allowed to oxidise  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$  in acidic medium, then  $\text{Fe}^{2+}$  oxidised will be :-  
 (A) more by  $\text{KMnO}_4$  (B) more by  $\text{K}_2\text{Cr}_2\text{O}_7$   
 (C) equal in both cases (D) can't be determined
17. If 10 g of  $\text{V}_2\text{O}_5$  is dissolved in acid and is reduced to  $\text{V}^{2+}$  by zinc metal, how many mole  $\text{I}_2$  could be reduced by the resulting solution if it is further oxidised to  $\text{VO}^{2+}$  ions ?  
 [Assume no change in state of  $\text{Zn}^{2+}$  ions] ( $V = 51, O = 16, I = 127$ ) :  
 (A) 0.11 mole of  $\text{I}_2$  (B) 0.22 mole of  $\text{I}_2$  (C) 0.055 mole of  $\text{I}_2$  (D) 0.44 mole of  $\text{I}_2$
18. Given that 50.0 mL of 0.01 M  $\text{Na}_2\text{S}_2\text{O}_3$  solution and  $5 \times 10^{-4}$  mole of  $\text{Cl}_2$  react according to equation,  

$$\text{Cl}_2(\text{g}) + \text{S}_2\text{O}_3^{2-} \longrightarrow \text{SO}_4^{2-} + \text{Cl}^- + \text{S}$$
  
 Answer the following :
- (i) The balanced molecular equation is :  
 (A)  $\text{Cl}_2 + \text{H}_2\text{O} + \text{Na}_2\text{S}_2\text{O}_3 \longrightarrow \text{Na}_2\text{SO}_4 + \text{S} + 2\text{HCl}$  (B)  $\text{Cl}_2 + \text{Na}_2\text{S}_2\text{O}_3 \longrightarrow 2\text{NaCl} + \text{Na}_2\text{SO}_4$   
 (C)  $\text{Cl}_2 + \text{S}_2\text{O}_3^{2-} \longrightarrow \text{SO}_4^{2-} + \text{S} + \text{Cl}^-$  (D) none of these
- (ii) How many moles of  $\text{S}_2\text{O}_3^{2-}$  are in the above sample :-  
 (A) 0.00050 (B) 0.0025 (C) 0.01 (D) 0.02
- (iii) How many equivalents of oxidising agents are in this sample for the above reaction :-  
 (A) 0.001 (B) 0.080 (C) 0.020 (D) 0.010
- (iv) What is the molarity of  $\text{Na}_2\text{SO}_4$  in this solution :-  
 (A) 0.080 M (B) 0.040 M (C) 0.020 M (D) 0.010 M
19. One gram of  $\text{Na}_3\text{AsO}_4$  is boiled with excess of solid KI in presence of strong HCl. The iodine evolved is absorbed in KI solution and titrated against 0.2 N hyposolution. Assuming the reaction to be  

$$\text{AsO}_4^{3-} + 2\text{H}^+ + 2\text{I}^- \longrightarrow \text{AsO}_3^{3-} + \text{H}_2\text{O} + \text{I}_2$$
  
 calculate the volume of thiosulphate hypo consumed. [Atomic weight of As = 75]  
 (A) 48.1 mL (B) 38.4 mL (C) 24.7 mL (D) 30.3 mL

## REDOX REACTION AND EQUIVALENT CONCEPT

20. Which of the following samples of reducing agents is/are chemically equivalent to 25 mL of 0.2 N  $\text{KMnO}_4$  to be reduced to  $\text{Mn}^{2+}$  and water :-  
 (A) 50 mL of 0.2 M  $\text{FeSO}_4$  to be oxidized to  $\text{Fe}^{3+}$       (B) 50 mL of 0.1 M  $\text{H}_3\text{AsO}_3$  to be oxidized to  $\text{H}_3\text{AsO}_4$   
 (C) 50 mL of 0.1 M  $\text{H}_2\text{O}_2$  to be oxidized to  $\text{H}^+$  and  $\text{O}_2$       (D) 25 mL of 0.1 M  $\text{SnCl}_2$  to be oxidized to  $\text{Sn}^{4+}$
21. Find the volume of strength of  $\text{H}_2\text{O}_2$  solution prepared by mixing of 250 mL of 3N  $\text{H}_2\text{O}_2$  & 750 mL of 1N  $\text{H}_2\text{O}_2$  solution  
 (A) 1.5 V      (B) 8.4 V      (C) 5.6 V      (D) 11.2 V
22. 0.3 g of an oxalate salt was dissolved in 100 mL solution. The solution required 90 mL of N/20  $\text{KMnO}_4$  for complete oxidation. The % of oxalate ion in salt is :-  
 (A) 33%      (B) 66%      (C) 70%      (D) 40%
23. A 0.518 g sample of limestone is dissolved in  $\text{HCl}$  and then the calcium is precipitated as  $\text{CaC}_2\text{O}_4$ . After filtering and washing the precipitate, it requires 40.0 mL of 0.250 N  $\text{KMnO}_4$  solution acidified with  $\text{H}_2\text{SO}_4$  to titrate it as. The percentage of  $\text{CaO}$  in the sample is :-  

$$\text{MnO}_4^- + \text{H}^+ + \text{C}_2\text{O}_4^{2-} \longrightarrow \text{Mn}^{2+} + \text{CO}_2 + 2\text{H}_2\text{O}$$
 (A) 54.0%      (B) 27.1%      (C) 42%      (D) 84%
24. In the reaction  $\text{CrO}_5 + \text{H}_2\text{SO}_4 \rightarrow \text{Cr}_2(\text{SO}_4)_3 + \text{H}_2\text{O} + \text{O}_2$  one mole of  $\text{CrO}_5$  will liberate how many moles of  $\text{O}_2$  :-  
 (A) 5/2      (B) 5/4      (C) 9/2      (D) 7/2
25. 25 mL of 0.50 M  $\text{H}_2\text{O}_2$  solution is added to 50 mL of 0.20 M  $\text{KMnO}_4$  in acid solution. Which of the following statement are true :-  
 (A) 0.010 mole of oxygen is liberated      (B) 0.005 mole of  $\text{KMnO}_4$  are left  
 (C) 0.030 g atom of oxygen gas is evolved      (D) 0.0025 mole  $\text{H}_2\text{O}_2$  does not react with  $\text{KMnO}_4$
26. Hydrogen peroxide in aqueous solution decomposes on warming to give oxygen according to the equation  

$$2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\ell) + \text{O}_2(\text{g})$$
 Under conditions where 1 mole of gas occupies 24  $\text{dm}^3$ . 100  $\text{cm}^3$  of XM solution of  $\text{H}_2\text{O}_2$  produces 3  $\text{dm}^3$  of  $\text{O}_2$ . Thus X is :-  
 (A) 2.5      (B) 1      (C) 0.5      (D) 0.25
27. Bottle (A) contain 320 mL of  $\text{H}_2\text{O}_2$  solution & labeled with 10 V  $\text{H}_2\text{O}_2$  & Bottle (B) contain 80 mL  $\text{H}_2\text{O}_2$  having normality 5N. If bottle (A) & bottle (B) mixed & solution filled in bottle (C). Select the correct label for bottle (C) in term of volume strength & in term of g / litre. :-  
 (A) 13.6 "V" & 41.285 g/L      (B) 11.2 "V" & 0.68 g/L  
 (C) 5.6 "V" & 0.68 g/L      (D) 5.6 "V" & 41.285 g/L
28. Temporary hardness is due to  $\text{HCO}_3^-$  of  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$ . It is removed by addition of  $\text{CaO}$ .  

$$\text{Ca}(\text{HCO}_3)_2 + \text{CaO} \rightarrow 2\text{CaCO}_3 + \text{H}_2\text{O}$$
 Mass of  $\text{CaO}$  required to precipitate 2 g  $\text{CaCO}_3$  is :-  
 (A) 2.00      (B) 0.56 g      (C) 0.28 g      (D) 1.12 g
29. An ore of iron, Wustite has the formula  $\text{Fe}_{0.93}\text{O}_{1.00}$ . The mole fraction of total iron present in the form of  $\text{Fe(II)}$  is :-  
 (A) 0.82      (B) 0.85      (C) 0.15      (D) 0.37
30.  $\text{HNO}_3$  oxidises  $\text{NH}_4^+$  ions to nitrogen and itself gets reduced to  $\text{NO}_2$ . The moles of  $\text{HNO}_3$  required by 1 mol of  $(\text{NH}_4)_2\text{SO}_4$  is :-  
 (A) 4      (B) 5      (C) 6      (D) 2
31. 25 mL of a 0.1 M solution of a stable cation of transition metal Z reacts exactly with 25 ml of 0.04 M acidified  $\text{KMnO}_4$  solution. Which of the following is most likely to represent the change in oxidation state of Z correctly :-  
 (A)  $\text{Z}^+ \rightarrow \text{Z}^{2+}$       (B)  $\text{Z}^{2+} \rightarrow \text{Z}^{3+}$       (C)  $\text{Z}^{3+} \rightarrow \text{Z}^{4+}$       (D)  $\text{Z}^{2+} \rightarrow \text{Z}^{4+}$

## CHEMISTRY FOR JEE MAIN & ADVANCED

32. 1 mol of iron (Fe) reacts completely with 0.65 mol  $O_2$  to give a mixture of only FeO and  $Fe_2O_3$ . Mole ratio of ferrous oxide to ferric oxide is :-  
 (A) 3 : 2 (B) 4 : 3 (C) 20 : 13 (D) none of these
33. The molar ratio of  $Fe^{++}$  to  $Fe^{+++}$  in a mixture of  $FeSO_4$  and  $Fe_2(SO_4)_3$  having equal number of sulphate ion in both ferrous and ferric sulphate is :-  
 (A) 1 : 2 (B) 3 : 2 (C) 2 : 3 (D) can't be determined
34. If a piece of iron gains 10% of its weight due to partial rusting into  $Fe_2O_3$ . The percentage of total iron that has rusted is :-  
 (A) 23 (B) 13 (C) 23.3 (D) 25.67
35. How many litres of  $Cl_2$  at S.T.P. will be liberated by oxidation of NaCl with 10 g  $KMnO_4$  :-  
 (A) 3.54 litres (B) 7.08 litres (C) 1.77 litres (D) none of these
36. During the disproportionation of iodine to iodide and iodate ions, the ratio of iodate and iodide ions formed in alkaline medium is :-  
 (A) 1 : 5 (B) 5 : 1 (C) 3 : 1 (D) 1 : 3
37.  $28 NO_3^- + 3 As_2S_3 + 4 H_2O \rightarrow 6 AsO_4^{3-} + 28 NO + 9 SO_4^{2-} + H^+$   
 What will be the equivalent mass of  $As_2S_3$  in above reaction  
 (A)  $\frac{M.wt.}{2}$  (B)  $\frac{M.wt.}{4}$  (C)  $\frac{M.wt.}{24}$  (D)  $\frac{M.wt.}{28}$
38.  $CN^-$  is oxidised by  $NO_3^-$  in presence of acid :  

$$a CN^- + b NO_3^- + c H^+ \longrightarrow (a + b) NO + a CO_2 + \frac{c}{2} H_2O$$
 What are the values of a, b, c in that order :  
 (A) 3,7,7 (B) 3,10,7 (C) 3,10,10 (D) 3,7,10
39. Which of the following solutions will exactly oxidize 25 mL of an acid solution of 0.1 M Fe (II) oxalate :-  
 (A) 25 mL of 0.1 M  $KMnO_4$  (B) 25 mL of 0.2 M  $KMnO_4$   
 (C) 25 mL of 0.6 M  $KMnO_4$  (D) 15 mL of 0.1 M  $KMnO_4$
40. 4.9 gm of  $K_2Cr_2O_7$  is taken to prepare 0.1 L of the solution. 10 mL of this solution is further taken to oxidise  $Sn^{2+}$  ion into  $Sn^{4+}$  ion  $Sn^{4+}$  so produced is used in 2<sup>nd</sup> reaction to prepare  $Fe^{3+}$  ion then the millimoles of  $Fe^{3+}$  ion formed will be (assume all other components are in sufficient amount) [Molar mass of  $K_2Cr_2O_7 = 294$  g].  
 (A) 5 (B) 20 (C) 10 (D) none of these
41. When ZnS is boiled with strong nitric acid, the products are zinc nitrate, sulphuric acid and nitrogen dioxide. What are the changes in the oxidation numbers of Zn, S and N.  
 (A) +2, +4, -1 (B) +2, +6, -2 (C) 0, +4, -2 (D) 0, +8, -1
42. When arsenic sulphide is boiled with NaOH, sodium arsenite and sodium thioarsenite are formed  
 $x As_2S_3 + y NaOH \longrightarrow Na_3AsO_3 + x Na_3AsS_3 + \frac{y}{2} H_2O$ . What are the values of x and y ?  
 (A) 1, 6 (B) 2, 8 (C) 2, 6 (D) 1, 4
43. An element forms two different sulphates in which its weight % is 28 and 37. What is the ratio of oxidation numbers of the element in these sulphates ?  
 (A) 1 : 2 (B) 1 : 3 (C) 2 : 1 (D) 3 : 2
44. The following equations are balanced atomwise and chargewise.  
 (i)  $Cr_2O_7^{2-} + 8H^+ + 3H_2O_2 \longrightarrow 2Cr^{3+} + 7H_2O + 3O_2$   
 (ii)  $Cr_2O_7^{2-} + 8H^+ + 5H_2O_2 \longrightarrow 2Cr^{3+} + 9H_2O + 4O_2$   
 (iii)  $Cr_2O_7^{2-} + 8H^+ + 7H_2O_2 \longrightarrow 2Cr^{3+} + 11H_2O + 5O_2$   
 The precise equation/equations representing the oxidation of  $H_2O_2$  is /are :  
 (A) (i) only (B) (ii) only (C) (iii) only (D) all the three



## REDOX REACTION AND EQUIVALENT CONCEPT

45. 35 mL sample of hydrogen peroxide gives of 500 mL of  $O_2$  at  $27^\circ C$  and 1 atm pressure. Volume strength of  $H_2O_2$  sample will be :-  
(A) 10 volume                      (B) 13 volumes                      (C) 11 volume                      (D) 12 volume
46. 20 mL of 0.1 M solution of compound  $Na_2CO_3 \cdot NaHCO_3 \cdot 2H_2O$  is titrated against 0.05 M HCl, x mL of HCl is used when phenolphthalein is used as an indicator and y mL of HCl is used when methyl orange is the indicator in two separate titrations. Hence  $(y - x)$  is :-  
(A) 40 mL                      (B) 80 mL                      (C) 120 mL                      (D) none of these
47. A mixture of  $H_2SO_4$  and  $H_2C_2O_4$  (oxalic acid) and some inert impurity weighing 3.185 g was dissolved in water and the solution made up to 1 litre, 10 mL of this solution required 3 mL of 0.1 N NaOH for complete neutralization. In another experiment 100 mL of the same solution in hot condition required 4 mL of 0.02M  $KMnO_4$  solution for complete reaction. The wt. % of  $H_2SO_4$  in the mixture was :-  
(A) 40                      (B) 50                      (C) 60                      (D) 80
48. 0.80 g of sample of impure potassium dichromate was dissolved in water and made upto 500 mL solution. 25 mL of this solution treated with excess of KI in acidic medium and  $I_2$  liberated required 24 mL of a sodium thiosulphate solution. 30 mL of this sodium thiosulphate solution required 15 mL of N/20 solution of pure potassium dichromate. What was the percentage of  $K_2Cr_2O_7$  in given sample?  
(A) 73.5 %                      (B) 75.3 %                      (C) 36.75 %                      (D) none of these
49. 0.10 g of a sample containing  $CuCO_3$  and some inert impurity was dissolved in dilute sulphuric acid and volume made up to 50 mL. This solution was added into 50 mL of 0.04 M KI solution where copper precipitates as  $CuI$  and  $I^-$  is oxidized into  $I_3^-$ . A 10 mL portion of this solution is taken for analysis, filtered and made up free  $I_3^-$  and then treated with excess of acidic permanganate solution. Liberated iodine required 20 mL of 2.5 mM sodium thiosulphate solution to reach the end point.  
Determine weight percentage of  $CuCO_3$  in the original sample.  
(A) 7.41                      (B) 74.1                      (C) 61.75                      (D) none of these
50. A 150 mL of solution of  $I_2$  is divided into two unequal parts. I part reacts with hypo solution in acidic medium. 15 mL of 0.4 M hypo was consumed. II part was added with 100 mL of 0.3 M NaOH solution. Residual base required 10 mL of 0.3 M  $H_2SO_4$  solution for complete neutralization. What was the initial concentration of  $I_2$  ?  
(A) 0.08 M                      (B) 0.1 M                      (C) 0.2 M                      (D) none of these

**Exercise # 2**

**Part # I**

**[Multiple Correct Choice Type Questions]**

- In the titration of  $K_2Cr_2O_7$  and ferrous sulphate, following data is obtained :  
 $V_1$  mL of  $K_2Cr_2O_7$  solution of molarity  $M_1$  requires  $V_2$  mL of  $FeSO_4$  solution of molarity  $M_2$ .  
 Which of the following relations is/are true for the above titration :  
 (A)  $6M_1V_1 = M_2V_2$       (B)  $M_1V_1 = 6M_2V_2$       (C)  $N_1V_1 = N_2V_2$       (D)  $M_1V_1 = M_2V_2$
- Choose the correct statement(s) :  
 (A) 1 mole of  $MnO_4^-$  ion can oxidise 5 moles of  $Fe^{2+}$  ion in acidic medium.  
 (B) 1 mole of  $Cr_2O_7^{2-}$  ion can oxidise 6 moles of  $Fe^{2+}$  ion in acidic medium.  
 (C) 1 mole of  $Cu_2S$  can be oxidised by 1.6 moles of  $MnO_4^-$  ion in acidic medium.  
 (D) 1 mole of  $Cu_2S$  can be oxidised by 1.33 moles of  $Cr_2O_7^{2-}$  ion in acidic medium.
- Which of the following samples of reducing agents is /are chemically equivalent to 25 mL of 0.2 N  $KMnO_4$  to be reduced to  $Mn^{2+}$  and water :  
 (A) 25 mL of 0.2 M  $FeSO_4$  to be oxidized to  $Fe^{3+}$   
 (B) 50 mL of 0.1 M  $H_3AsO_3$  to be oxidized to  $H_3AsO_4$   
 (C) 25 mL of 0.1 M  $H_2O_2$  to be oxidized to  $H^+$  and  $O_2$   
 (D) 25 mL of 0.1 M  $SnCl_2$  to be oxidized to  $Sn^{4+}$
- 0.1 M solution of KI reacts with excess of  $H_2SO_4$  and  $KIO_3$  solutions, according to equation  
 $5I^- + IO_3^- + 6H^+ \longrightarrow 3I_2 + 3H_2O$ ; which of the following statements is/are correct :  
 (A) 200 mL of the KI solution react with 0.004 mole  $KIO_3$ .  
 (B) 100 mL of the KI solution reacts with 0.006 mole of  $H_2SO_4$ .  
 (C) 0.5 litre of the KI solution produced 0.005 mole of  $I_2$ .  
 (D) Equivalent weight of  $KIO_3$  is equal to  $\left(\frac{\text{Molecular Weight}}{5}\right)$ .
- Consider the redox reaction  $2S_2O_3^{2-} + I_2 \longrightarrow S_4O_6^{2-} + 2I^-$  :  
 (A)  $S_2O_3^{2-}$  gets reduced to  $S_4O_6^{2-}$       (B)  $S_2O_3^{2-}$  gets oxidised to  $S_4O_6^{2-}$   
 (C)  $I_2$  gets reduced to  $I^-$       (D)  $I_2$  gets oxidised to  $I^-$
- There are two sample of HCl having molarity 1N and 0.25 N. Find volume of these sample taken in order to prepare 0.75 N HCl solution. (Assume no water is used) :  
 (A) 20 mL, 10 mL      (B) 100 mL, 50 mL      (C) 40 mL, 20 mL      (D) 50 mL, 25 mL
- To a 25 ml  $H_2O_2$  solution excess acidified solution of KI was added. The iodine liberated 20 ml of 0.3 N sodium thiosulphate solution. Use these data to choose the correct statements from the following :  
 (A) The weight of  $H_2O_2$  present in 25 ml solution is 0.102 g  
 (B) The molarity of  $H_2O_2$  solution is 0.12 M  
 (C) The weight of  $H_2O_2$  present in 1 L of the solution is 0.816 g  
 (D) The volume strength of  $H_2O_2$  is 1.344 L



16. The molality of a sulphuric acid solution is 0.2. Calculate the total weight of the solution having 1000 gm of solvent.  
 (A) 1000 g (B) 1098.6 g (C) 980.4 g (D) 1019.6g
17. If  $\text{H}_2\text{SO}_4$  is formed from its elements by taking  $6.023 \times 10^{23}$  atom of 'O' 5.6 litre of  $\text{H}_2$  gas at STP and 8 gm S then  
 (A) 0.125 moles of  $\text{H}_2\text{SO}_4$  are formed (B) 0.25 moles of  $\text{H}_2\text{SO}_4$  are formed  
 (C) no moles of 'S' are left (D) 1/4 mole of  $\text{O}_2$  is left
18. A 5L vessel contains 2.8 g of  $\text{N}_2$ . When heated to 1800 K, 30% molecules are dissociated into atoms.  
 (A) Total no. of moles in the container will be 0.13  
 (B) Total no. of molecules in the container will be close to  $0.421 \times 10^{23}$ .  
 (C) Total no. of moles in the container will be 0.098.  
 (D) All of these are correct.
19. The density of air is  $0.001293 \text{ g/cm}^3$  at STP. Identify which of the following statement is correct  
 (A) Vapour density is 14.48  
 (B) Molecular weight is 28.96  
 (C) Vapour density is  $0.001293 \text{ g/cm}^3$   
 (D) Vapour density and molecular weight cannot be determined.
20. Which of the following has same mass  
 (A) 1.0 mole of  $\text{O}_2$  (B)  $3.01 \times 10^{23}$  molecules of  $\text{SO}_2$   
 (C) 0.5 moles of  $\text{CO}_2$  (D) 1 g atom of sulphur

Part # II

[Assertion & Reason Type Questions]

These questions contains, Statement I (assertion) and Statement II (reason).

- (A) Statement-I is true, Statement-II is true ; Statement-II is correct explanation for Statement-I.  
 (B) Statement-I is true, Statement-II is true ; Statement-II is NOT a correct explanation for statement-I  
 (C) Statement-I is true, Statement-II is false  
 (D) Statement-I is false, Statement-II is true

1. **Statement-I** : Oxidation involves loss of electrons and reduction involves gain of electrons.  
**Statement-II** : The overall reaction in which oxidation and reduction occur simultaneously is called redox reaction.
2. **Statement-I** : The oxidation state of superoxide ion in  $\text{KO}_2$ ,  $\text{CsO}_2$  and  $\text{RbO}_2$  is  $-1/2$ .  
**Statement-II** : Since the oxidation state of an alkali metal in any compound is always +1, the oxidation state of oxygen is  $-1/2$  in the  $\text{O}_2^-$  ion.
3. **Statement-I** : Among  $\text{Br}^-$ ,  $\text{O}_2^{2-}$ ,  $\text{H}^-$  and  $\text{NO}_3^-$ , the ions that could not act as oxidising agents are  $\text{Br}^-$  and  $\text{H}^-$ .  
**Statement-II** :  $\text{Br}^0$  and  $\text{H}^-$  could not be reduced.
4. **Statement-I** :  $\text{H}_2\text{SO}_4$  cannot act as reducing agent.  
**Statement-II** : Sulphur cannot increase its oxidation number beyond +6.

5. **Statement-I :** In the redox reaction  

$$8 \text{H}^+ (\text{aq}) + 4 \text{NO}_3^- + 6 \text{Cl}^- + \text{Sn} (\text{s}) \rightarrow \text{SnCl}_6^{2-} + 4 \text{NO}_2 + 4 \text{H}_2\text{O}$$
 the reducing agent is Sn (s),  
**Statement-II :** In balancing half reaction,  $\text{S}_2\text{O}_3^{2-} \rightarrow \text{S}(\text{s})$ , the number of electrons added on the left is 4.
6. **Statement-1 :** The molality and molarity of very dilute aqueous solutions differ very little.  
**Statement-2 :** The density of water is about  $1.0 \text{ g cm}^{-3}$  at room temperature.
7. **Statement-1 :** For calculating the molality or the mole fraction of solute, if the molarity is known, it is necessary to know the density of the solution.  
**Statement-2 :** Molality, molarity and the mole fraction of solute can be calculated from the weight percentage and the density of the solution
8. **Statement-1 :** Laboratory reagents are usually made up to a specific molarity rather than a given molality.  
**Statement-2 :** The volume of a liquid is more easily measured than its mass.
9. **Statement-1 :** Molality and mole fraction concentration units do not change with temperature.  
**Statement-2 :** These units are not defined in terms of any volume.
10. **Statement-1 :** A one molal solution prepared at  $20^\circ\text{C}$  will retain the same molality at  $100^\circ\text{C}$ , provided there is no loss of solute or solvent on heating.  
**Statement-2 :** Molality is independent of temperature.
11. **Statement-1 :** In the reaction,  $\text{MnO}_4^- + 5\text{Fe}^{2+} + 8\text{H}^+ \longrightarrow \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O}$ ,  $\text{MnO}_4^-$  acts as oxidising agent.  
**Statement-2 :** In the above reaction, n-factor for  $\text{MnO}_4^-$  is 5.
12. **Statement-1 :** If 200 mL of 0.1N NaOH is added to 200 mL of 0.1N  $\text{H}_2\text{SO}_4$  solution, then the resulting solution is acidic.  
**Statement-2 :** If milliequivalents of acid is greater than milliequivalents of base, then upon mixing the solution is acidic.
13. **Statement-1 :** Equivalent weight of  $\text{FeC}_2\text{O}_4$  in the reaction,  

$$\text{FeC}_2\text{O}_4 + \text{Oxidising agent} \longrightarrow \text{Fe}^{3+} + \text{CO}_2$$
 is  $M/3$ , where M is molar mass of  $\text{FeC}_2\text{O}_4$ .  
**Statement-2 :** In the above reaction, total two moles of electrons are given up by 1 mole of  $\text{FeC}_2\text{O}_4$  to the oxidising agent.

## Exercise # 3

Part # I

[Matrix Match Type Questions]

1. **Column-I**
- (A) Eq. wt. =  $\frac{\text{Molecular weight}}{33}$
- (B) Eq. wt. =  $\frac{\text{Molecular weight}}{27}$
- (C) Eq. wt. =  $\frac{\text{Molecular weight}}{28}$
- (D) Eq. wt. =  $\frac{\text{Molecular weight}}{24}$
- Column-II**
- (p) When  $\text{CrI}_3$  oxidises into  $\text{Cr}_2\text{O}_7^{2-}$  and  $\text{IO}_4^-$
- (q) When  $\text{Fe}(\text{SCN})_2$  oxidises into  $\text{Fe}^{3+}$ ,  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$  and  $\text{NO}_3^-$
- (r) When  $\text{NH}_4\text{SCN}$  oxidizes into  $\text{SO}_4^{2-}$ ,  $\text{CO}_3^{2-}$  and  $\text{NO}_3^-$
- (s) When  $\text{As}_2\text{S}_3$  oxidises into  $\text{AsO}_3^-$  and  $\text{SO}_4^{2-}$
2. **Column-I**
- (A) When  $\text{Bi}_2\text{S}_3$  converted into  $\text{Bi}^{5+}$  and S
- (B) When  $\text{Al}_2(\text{Cr}_2\text{O}_7)_3$  reduced into  $\text{Cr}^{3+}$  in acidic medium
- (C) When  $\text{FeS}_2$  converted into  $\text{Fe}_2\text{O}_3$  and  $\text{SO}_2$
- (D) When  $\text{Mn}(\text{NO}_3)_2$  converted into  $\text{MnO}_4^{2-}$  and NO
- Column-II**
- (p) 18
- (q) 11
- (r) 2
- (s) 10
3. **Column-I**
- (A)  $\text{P}_2\text{H}_4 \longrightarrow \text{PH}_3 + \text{P}_4\text{H}_2$
- (B)  $\text{I}_2 \longrightarrow \text{I}^- + \text{IO}_3^-$
- (C)  $\text{MnO}_4^- + \text{Mn}^{2+} + \text{H}_2\text{O} \longrightarrow \text{Mn}_3\text{O}_4 + \text{H}^+$
- (D)  $\text{H}_3\text{PO}_2 \longrightarrow \text{PH}_3 + \text{H}_3\text{PO}_3$
- Column-II**
- (p)  $E = \frac{3M}{4}$
- (q)  $E = \frac{3M}{5}$
- (r)  $E = \frac{15M}{26}$
- (s)  $E = \frac{5M}{6}$
4. **Column-I**
- (A) Molarity
- (B) Molality
- (C) Mole fraction
- (D) Mass %
- Column-II**
- (p) Dependent on temperature
- (q)  $\frac{M_A \times n_A}{n_A M_A + n_B M_B} \times 100$
- (r) Independent of temperature
- (s)  $\frac{X_A}{X_B M_B} \times 1000$

Where  $M_A$ ,  $M_B$  are molar masses,  $n_A$ ,  $n_B$  are no of moles &  $X_A$ ,  $X_B$  are mole fractions of solute and solvent respectively.

## REDOX REACTION AND EQUIVALENT CONCEPT


5. **Column-I**
- (A) 100 ml of 0.2 M  $\text{AlCl}_3$  solution + 400 ml
- (B) 50 ml of 0.4 M  $\text{KCl}$  + 50 ml  $\text{H}_2\text{O}$
- (C) 30 ml of 0.2 M  $\text{K}_2\text{SO}_4$  + 70 ml  $\text{H}_2\text{O}$
- (D) 200 ml 24.5% (w/v)  $\text{H}_2\text{SO}_4$
- Column-II**
- (p) Total concentration of cation(s) = 0.12 M of 0.1 M  $\text{HCl}$  solution
- (q)  $[\text{SO}_4^{2-}] = 0.06 \text{ M}$
- (r)  $[\text{SO}_4^{2-}] = 2.5 \text{ M}$
- (s)  $[\text{Cl}^-] = 0.2 \text{ M}$
6. **Column-I**
- (A) 4.1 g  $\text{H}_2\text{SO}_3$
- (B) 4.9 g  $\text{H}_3\text{PO}_4$
- (C) 4.5 g oxalic acid ( $\text{H}_2\text{C}_2\text{O}_4$ )
- (D) 5.3 g  $\text{Na}_2\text{CO}_3$
- Column-II**
- (p) 200 mL of 0.5 N base is used for complete neutralization
- (q) 200 millimoles of oxygen atoms
- (r) Central atom is in its highest oxidation number
- (s) May react with an oxidising agent
7. **Column-I**
- (A)  $\text{Sn}^{+2} + \text{MnO}_4^-$  (acidic)  
3.5 mole    1.2 mole
- (B)  $\text{H}_2\text{C}_2\text{O}_4 + \text{MnO}_4^-$  (acidic)  
8.4 mole    3.6 mole
- (C)  $\text{S}_2\text{O}_3^{2-} + \text{I}_2$   
7.2 mole    3.6 mole
- (D)  $\text{Fe}^{+2} + \text{Cr}_2\text{O}_7^{2-}$  (acidic)  
9.2 mole    1.6 mole
- Column-II**
- (p) Amount of oxidant available decides the number of electrons transfer
- (q) Amount of reductant available decides the number of electrons transfer
- (r) Number of electrons involved per mole of oxidant > Number of electrons involved per mole of reductant
- (s) Number of electrons involved per mole of oxidant < Number of electrons involved per mole of reductant.

### Part # II

### [Comprehension Type Questions]

#### Comprehension # 1

The strength of  $\text{H}_2\text{O}_2$  is expressed in several ways like molarity, normality, % (w/V), volume strength, etc. The strength of "10 V" means 1 volume of  $\text{H}_2\text{O}_2$  on decomposition gives 10 volumes of oxygen at STP or 1 litre of  $\text{H}_2\text{O}_2$  gives 10 litre of  $\text{O}_2$  at STP. The decomposition of  $\text{H}_2\text{O}_2$  is shown as under :



$\text{H}_2\text{O}_2$  can acts as oxidising as well as reducing agent, as oxidizing agent  $\text{H}_2\text{O}_2$  converted into  $\text{H}_2\text{O}$  and as reducing agent  $\text{H}_2\text{O}_2$  converted into  $\text{O}_2$ , both cases it's n-factor is 2.

$\therefore$  Normality of  $\text{H}_2\text{O}_2$  solution =  $2 \times$  Molarity of  $\text{H}_2\text{O}_2$  solution

1. What is the molarity of "11.2 V" of  $\text{H}_2\text{O}_2$  ?
- (A) 1 M                                      (B) 2 M                                      (C) 5.6 M                                      (D) 11.2 M
2. What is the percentage strength (% w/V) of "11.2 V"  $\text{H}_2\text{O}_2$  ?
- (A) 1.7                                      (B) 3.4                                      (C) 34                                      (D) none of these
3. 20 mL of  $\text{H}_2\text{O}_2$  solution is reacted with 80 mL of 0.05 M  $\text{KMnO}_4$  in acidic medium then what is the volume strength of  $\text{H}_2\text{O}_2$  ?
- (A) 2.8                                      (B) 5.6                                      (C) 11.2                                      (D) none of these

4. 40 g  $\text{Ba}(\text{MnO}_4)_2$  (mol. wt. = 375) sample containing some inert impurities in acidic medium is completely reacted with 125 mL of "33.6 V" of  $\text{H}_2\text{O}_2$ . What is the percentage purity of the sample?  
 (A) 28.12% (B) 70.31% (C) 85% (D) none of these

### Comprehension # 2

Oleum is considered as a solution of  $\text{SO}_3$  in  $\text{H}_2\text{SO}_4$ , which is obtained by passing  $\text{SO}_3$  in solution of  $\text{H}_2\text{SO}_4$ . When 100 g sample of oleum is diluted with desired weight of  $\text{H}_2\text{O}$  then the total mass of  $\text{H}_2\text{SO}_4$  obtained after dilution is known as % labelling in oleum.

For example, a oleum bottle labelled as '109%  $\text{H}_2\text{SO}_4$ ' means the 109 g total mass of pure  $\text{H}_2\text{SO}_4$  will be formed when 100 g of oleum is diluted by 9 g of  $\text{H}_2\text{O}$  which combines with all the free  $\text{SO}_3$  present in oleum to form  $\text{H}_2\text{SO}_4$  as  $\text{SO}_3 + \text{H}_2\text{O} \longrightarrow \text{H}_2\text{SO}_4$

- What is the % of free  $\text{SO}_3$  in an oleum that is labelled as '104.5 %  $\text{H}_2\text{SO}_4$ '?  
 (A) 10 (B) 20 (C) 40 (D) none of these
- 9.0 g water is added into oleum sample labelled as '112%  $\text{H}_2\text{SO}_4$ ' then the amount of free  $\text{SO}_3$  remaining in the solution is :  
 (A) 14.93 L at STP (B) 7.46 L at STP (C) 3.73 L at STP (D) 11.2 L at STP
- If excess water is added into a bottle sample labelled as '112 %  $\text{H}_2\text{SO}_4$ ' and is reacted with 5.3 g  $\text{Na}_2\text{CO}_3$ , then find the volume of  $\text{CO}_2$  evolved at 1 atm pressure and 300 K temperature after the completion of the reaction :  
 (A) 2.46 L (B) 24.6 L (C) 1.23 L (D) 12.3 L
- 1 g of oleum sample is diluted with water. The solution required 54 mL of 0.4 N NaOH for complete neutralization. The % of free  $\text{SO}_3$  in the sample is :  
 (A) 74 (B) 26 (C) 20 (D) none of these

### Comprehension # 3


$$\text{Equivalent weight} = \frac{\text{Molecular weight} / \text{Atomic weight}}{n - \text{factor}}$$

n-factor is very important in redox as well as non-redox reactions. With the help of n-factor we can predicts the molar ratio of the reactant species taking part in reactions. The reciprocal of n-factor's ratio of the reactants is the molar ratio of the reactants.

In general n-factor of acid/base is number of moles of  $\text{H}^+$  /  $\text{OH}^-$  furnished per mole of acid/base. n-factor of a reactant is no. of moles of electrons lost or gained per mole of reactant.

#### Example 1

- In acidic medium :  $\text{KMnO}_4 (n = 5) \longrightarrow \text{Mn}^{2+}$
- In basic medium :  $\text{KMnO}_4 (n = 3) \longrightarrow \text{Mn}^{2+}$
- In neutral medium :  $\text{KMnO}_4 (n = 1) \longrightarrow \text{Mn}^{6+}$



Total no. of moles of  $e^-$  lost by 1 mole of  $\text{FeC}_2\text{O}_4 = 1 + 1 \times 2 \Rightarrow 3$

$\therefore$  n-factor of  $\text{FeC}_2\text{O}_4 = 3$

- n-factor of  $\text{Ba}(\text{MnO}_4)_2$  in acidic medium is :  
 (A) 2 (B) 6 (C) 10 (D) none of these



## REDOX REACTION AND EQUIVALENT CONCEPT

2. For the reaction,
- $$\text{H}_3\text{PO}_2 + \text{NaOH} \longrightarrow \text{NaH}_2\text{PO}_2 + \text{H}_2\text{O}$$
- What is the equivalent weight of  $\text{H}_3\text{PO}_2$ ? (mol. wt. is M)
- (A) M                      (B) M/2                      (C) M/3                      (D) none of these
3. For the reaction,  $\text{Fe}_{0.95}\text{O}$  (molar mass : M)  $\longrightarrow$   $\text{Fe}_2\text{O}_3$ . What is the eq. wt. of  $\text{Fe}_{0.95}\text{O}$ ?
- (A)  $\frac{M}{0.85}$                       (B)  $\frac{M}{0.95}$                       (C)  $\frac{M}{0.8075}$                       (D) none of these
4. In the reaction,  $x\text{VO} + y\text{Fe}_2\text{O}_3 \longrightarrow \text{FeO} + \text{V}_2\text{O}_5$ . What is the value of x and y respectively?
- (A) 1, 1                      (B) 2, 3                      (C) 3, 2                      (D) none of these

### Comprehension # 4

Some amount of "20V"  $\text{H}_2\text{O}_2$  is mixed with excess of acidified solution of KI. The iodine so liberated required 200 mL of 0.1 N  $\text{Na}_2\text{S}_2\text{O}_3$  for titration.

1. The volume of  $\text{H}_2\text{O}_2$  solution is :
- (A) 11.2 mL                      (B) 37.2 mL                      (C) 5.6 mL                      (D) 22.4 mL
2. The mass of  $\text{K}_2\text{Cr}_2\text{O}_7$  needed to oxidise the above volume of  $\text{H}_2\text{O}_2$  solution is :
- (A) 3.6 g                      (B) 0.8 g                      (C) 4.2 g                      (D) 0.98 g
3. The volume of  $\text{O}_2$  at STP that would be liberated by above  $\text{H}_2\text{O}_2$  solution is :
- (A) 56 mL                      (B) 112 mL                      (C) 168 mL                      (D) 224 mL

### Comprehension # 5

#### Equivalent Mass

The equivalent mass of a substance is defined as the number of parts by mass of it which combine with or displace 1.0078 parts by mass of hydrogen, 8 parts by mass of oxygen and 35.5 parts by mass of chlorine.

The equivalent mass of a substance expressed in grams is called **gram equivalent mass**.

The equivalent mass of a substance is not constant. It depends upon the reaction in which the substance is participating. A compound may have different equivalent mass in different chemical reactions and under different experimental conditions.

#### (A) Equivalent mass of an acid

It is the mass of an acid in grams which contains 1.0078 g of replaceable  $\text{H}^+$  ions or it is the mass of acid which contains one mole of replaceable  $\text{H}^+$  ions. It may be calculated as :

$$\text{Equivalent mass of acid} = \frac{\text{Molecular mass of acid}}{\text{Basicity of acid}}$$

Basicity of acid = number of replaceable hydrogen atoms present in one molecule of acid

#### (B) Equivalent mass of a base

It is the mass of the base which contains one mole of replaceable  $\text{OH}^-$  ions in molecule.

$$\text{Equivalent mass of base} = \frac{\text{Molecular mass of base}}{\text{Acidity of base}}$$

Acidity of base = Number of replaceable  $\text{OH}^-$  ions present in one molecule of the base

**Equivalent mass of an oxidising agent**

(a) **Electron concept** : Equivalent mass of oxidising agent =  $\frac{\text{Molecular mass of oxidising agent}}{\text{Number of electrons gained by one molecule}}$

(b) **Oxidation number concept** : Equivalent mass of oxidising agent =  $\frac{\text{Molecular mass of oxidising agent}}{\text{Total change in oxidation number per molecule of oxidising agent}}$

- Equivalent mass of  $\text{Ba}(\text{MnO}_4)_2$  in acidic medium is : (where M stands for molar mass)  
 (A)  $M/5$                       (B)  $M/6$                       (C)  $M/10$                       (D)  $M/2$
- Equivalent mass of  $\text{Fe}_{0.9}\text{O}$  in reaction with acidic  $\text{K}_2\text{Cr}_2\text{O}_7$  is : (M = Molar mass)  
 (A)  $7M/10$                       (B)  $10M/7$                       (C)  $7M/9$                       (D)  $9M/7$
- Equivalent weight of oxalic acid salt in following reaction is : (Atomic masses : O = 16, C = 12, K = 39)  
 $\text{H}_2\text{C}_2\text{O}_4 + \text{Ca}(\text{OH})_2 \longrightarrow \text{CaC}_2\text{O}_4 + \text{H}_2\text{O}$   
 (A) 90                      (B) 45                      (C) 64                      (D) 128

## Exercise # 4

## [Subjective Type Questions]

- $\text{KMnO}_4$  oxidizes  $\text{X}^{n+}$  ion to  $\text{XO}_3^-$ , itself changing to  $\text{Mn}^{2+}$  in acid medium.  $2.68 \times 10^{-3}$  mole of  $\text{X}^{n+}$  requires  $1.61 \times 10^{-3}$  mole of  $\text{MnO}_4^-$ . What is the value of  $n$ ? Also calculate the atomic mass of  $\text{X}$ , if the weight of 1 g equivalent of  $\text{XCl}_n$  is 56.
- In a quantitative determination of iron in an ore, an analyst converted 0.40 g, of the ore into its ferrous. This required 40.00 mL of 0.1 N solution of  $\text{KMnO}_4$  for titration.

  - How many milliequivalents of  $\text{KMnO}_4$  does 40.00 mL of 0.1 N solution represent?
  - How many equivalents of iron were present in the sample of the ore taken for analysis?
  - How many grams of iron were present in the sample?
  - What is the percentage of iron in the ore?
  - What is the molarity of  $\text{KMnO}_4$  solution used?
  - How many moles of  $\text{KMnO}_4$  were used for titration? ( $\text{Fe} = 56$ )
- The mixture of  $\text{CuS}$  (molar weight =  $M_1$ ) and  $\text{Cu}_2\text{S}$  (molecular weight =  $M_2$ ) oxidised by  $\text{KMnO}_4$  (molecular weight =  $M_3$ ) in acidic medium, the product obtained are  $\text{Cu}^{2+}$ ,  $\text{SO}_2$ . Find the equivalent weight of  $\text{CuS}$ ,  $\text{Cu}_2\text{S}$  and  $\text{KMnO}_4$  respectively.
- Calculate the oxidation number of underlined elements in the following compounds :

(a) $\text{K}[\underline{\text{C}}(\text{C}_2\text{O}_4)_2(\text{NH}_3)_2]$	(b) $\text{K}_4\underline{\text{P}}_2\text{O}_7$	(c) $\underline{\text{C}}\text{rO}_2\text{Cl}_2$
(d) $\text{Na}_2[\underline{\text{F}}\text{e}(\text{CN})_5\text{NO}^+]$	(e) $\underline{\text{Mn}}_3\text{O}_4$	(f) $\text{Ca}(\underline{\text{C}}\text{lO}_2)_2$
(g) $[\underline{\text{F}}\text{e}(\text{NO})(\text{H}_2\text{O})_5]\text{SO}_4$	(h) $\underline{\text{Zn}}\text{O}_2^{2-}$	(c) $\underline{\text{F}}\text{e}_{0.93}\text{O}$
- Write balanced net ionic equation for the following reactions in acidic solution.

  - $\text{S}_4\text{O}_6^{2-}(\text{aq}) + \text{Al}(\text{s}) \longrightarrow \text{H}_2\text{S}(\text{aq}) + \text{Al}^{3+}(\text{aq})$
  - $\text{S}_2\text{O}_3^{2-}(\text{aq}) + \text{Cr}_2\text{O}_7^{2-}(\text{aq}) \longrightarrow \text{S}_4\text{O}_6^{2-}(\text{aq}) + \text{Cr}^{3+}(\text{aq})$
  - $\text{ClO}_3^-(\text{aq}) + \text{As}_2\text{S}_3(\text{s}) \longrightarrow \text{Cl}^-(\text{aq}) + \text{H}_2\text{AsO}_4^-(\text{aq}) + \text{HSO}_4^-(\text{aq})$
  - $\text{IO}_3^-(\text{aq}) + \text{Re}(\text{s}) \longrightarrow \text{ReO}_4^-(\text{aq}) + \text{I}^-(\text{aq})$
  - $\text{HSO}_4^-(\text{aq}) + \text{As}_4(\text{s}) + \text{Pb}_3\text{O}_4(\text{s}) \longrightarrow \text{PbSO}_4(\text{s}) + \text{H}_2\text{AsO}_4^-(\text{aq})$
  - $\text{HNO}_2(\text{aq}) \longrightarrow \text{NO}_3^- + \text{NO}(\text{g})$
- Write balanced net ionic equations for the following reactions in basic solution :

  - $\text{C}_4\text{H}_4\text{O}_6^{2-}(\text{aq}) + \text{ClO}_3^-(\text{aq}) \longrightarrow \text{CO}_3^{2-}(\text{aq}) + \text{Cl}^-(\text{aq})$
  - $\text{Al}(\text{s}) + \text{BiONO}_3(\text{s}) \longrightarrow \text{Bi}(\text{s}) + \text{NH}_3(\text{aq}) + \text{Al}(\text{OH})_4^-(\text{aq})$
  - $\text{H}_2\text{O}_2(\text{aq}) + \text{Cl}_2\text{O}_7(\text{aq}) \longrightarrow \text{ClO}_2^-(\text{aq}) + \text{O}_2(\text{g})$
  - $\text{Ti}_2\text{O}_3(\text{s}) + \text{NH}_2\text{OH}(\text{aq}) \longrightarrow \text{TiOH}(\text{s}) + \text{N}_2(\text{g})$
  - $\text{Cu}(\text{NH}_3)_4^{2+}(\text{aq}) + \text{S}_2\text{O}_4^{2-}(\text{aq}) \longrightarrow \text{SO}_3^{2-}(\text{aq}) + \text{Cu}(\text{s}) + \text{NH}_3(\text{aq})$
  - $\text{Mn}(\text{OH})_2(\text{s}) + \text{MnO}_4^-(\text{aq}) \longrightarrow \text{MnO}_2(\text{s})$

## CHEMISTRY FOR JEE MAIN & ADVANCED

7. Consider the reaction  $\text{H}^+ + \text{IO}_4^- + \text{I}^- \rightarrow \text{I}_2 + \text{H}_2\text{O}$ . Find the ratio of coefficients of  $\text{IO}_4^-$ ,  $\text{I}^-$  and  $\text{I}_2$ .
8. A dilute solution of  $\text{H}_2\text{SO}_4$  is made by adding 5 mL of 3N  $\text{H}_2\text{SO}_4$  to 245 mL of water. Find the normality and molarity of the solution.
9. What volume at NTP of gaseous ammonia will be required to be passed into 30 cc of N –  $\text{H}_2\text{SO}_4$  solution to bring down the acid strength of the latter to 0.2 N.
10. 10 g  $\text{CaCO}_3$  were dissolved in 250 mL of M HCl and the solution was boiled. What volume of 2 M KOH would be required to equivalence point after boiling? Assume no change in volume during boiling.
11.  $\text{H}_3\text{PO}_4$  is a tri basic acid and one of its salt is  $\text{NaH}_2\text{PO}_4$ . What volume of 1 M NaOH solution should be added to 12 g of  $\text{NaH}_2\text{PO}_4$  to convert it into  $\text{Na}_3\text{PO}_4$ ?
12. 1.64 g of mixture of  $\text{CaCO}_3$  and  $\text{MgCO}_3$  was dissolved in 50 mL of 0.8 M HCl. The excess of acid required 16 mL of 0.25 M NaOH for neutralization. Calculate the percentage of  $\text{CaCO}_3$  and  $\text{MgCO}_3$  in the sample.
13. A solution containing 4.2 g of KOH and  $\text{Ca}(\text{OH})_2$  is neutralized by an acid. It consumes 0.1 equivalent of acid, calculate the percentage composition of the sample.
14. How many mL of 0.1 N HCl are required to react completely with 1 g mixture of  $\text{Na}_2\text{CO}_3$  and  $\text{NaHCO}_3$  containing equimolar amounts of two?
15. 0.5 g of fuming  $\text{H}_2\text{SO}_4$  (oleum) is diluted with water. The solution requires 26.7 mL of 0.4N NaOH for complete neutralization. Find the % of free  $\text{SO}_3$  in the sample of oleum.
16. 1.5 g of chalk were treated with 10 mL of 4N – HCl. The chalk was dissolved and the solution made to 100 mL, 25 mL of this solution required 18.75 mL of 0.2 N – NaOH solution for complete neutralisation. Calculate the percentage of pure  $\text{CaCO}_3$  in the sample of chalk?
17. A solution contains  $\text{Na}_2\text{CO}_3$  and  $\text{NaHCO}_3$ . 20 mL of this solution required 4 mL of 1N – HCl for titration with Ph indicator. The titration was repeated with the same volume of the solution but with MeOH. 10.5 mL of 1 – N HCl was required this time. Calculate the amount of  $\text{Na}_2\text{CO}_3$  &  $\text{NaHCO}_3$ .
18. A solution contains a mix of  $\text{Na}_2\text{CO}_3$  and NaOH. Using Ph as indicator 25 mL of mixture required 19.5 mL of 0.995 N HCl for the end point. With MeOH, 25 mL of the solution required 25 mL of the same HCl for the end point. Calculate g/L of each substance in the mixture.
19. It required 40.05 mL of 1 M  $\text{Ce}^{4+}$  to titrate 20 mL of 1 M  $\text{Sn}^{2+}$  to  $\text{Sn}^{4+}$ . What is the oxidation state of the cerium in the product.
20. A volume of 12.53 mL of 0.05093 M  $\text{SeO}_2$  reacted with exactly 25.52 mL of 0.1 M  $\text{CrSO}_4$ . In the reaction,  $\text{Cr}^{2+}$  was oxidized to  $\text{Cr}^{3+}$ . To what oxidation state was selenium converted by the reaction.
21. Pottassium acid oxalate  $\text{K}_2\text{C}_2\text{O}_4 \cdot 3\text{H}_2\text{C}_2\text{O}_4 \cdot 4\text{H}_2\text{O}$  can be oxidized by  $\text{MnO}_4^-$  in acid medium. Calculate the volume of 0.1 M  $\text{KMnO}_4$  reacting in acid solution with one gram of the acid oxalate.
22. 200 mL of a solution of mixture of NaOH and  $\text{Na}_2\text{CO}_3$  was first titrated with Ph and  $\frac{\text{N}}{10}$  HCl. 17.5 mL of HCl was required for end point. After this MeOH was added and 2.5 mL of same HCl was again required for next end point. Find out amounts of NaOH and  $\text{Na}_2\text{CO}_3$  in the mix.

## REDOX REACTION AND EQUIVALENT CONCEPT

23. A solution contains  $\text{Na}_2\text{CO}_3$  and  $\text{NaHCO}_3$ . 10 mL of this requires 2 mL of 0.1 M  $\text{H}_2\text{SO}_4$  for neutralisation using Ph indicator. MeOH is then added when a further 2.5 mL of 0.2 M  $\text{H}_2\text{SO}_4$  was needed. Calculate strength of  $\text{Na}_2\text{CO}_3$  and  $\text{NaHCO}_3$ .
24. A sample containing  $\text{Na}_2\text{CO}_3$  &  $\text{NaOH}$  is dissolved in 100 mL solution. 10 mL of this solution requires 25 mL of 0.1 N HCl when Ph is used as indicator. If MeOH is used as indicator 10 mL of same solution requires 30 mL of same HCl. Calculate % of  $\text{Na}_2\text{CO}_3$  and  $\text{NaOH}$  in the sample.
25. A 1.0 g sample of  $\text{H}_2\text{O}_2$  solution containing x%  $\text{H}_2\text{O}_2$  by mass requires  $x \text{ cm}^3$  of a  $\text{KMnO}_4$  solution for complete oxidation under acidic conditions. Calculate the normality of  $\text{KMnO}_4$  solution.
26. Metallic tin in the presence of HCl is oxidized by  $\text{K}_2\text{Cr}_2\text{O}_7$  to stannic chloride,  $\text{SnCl}_4$ . What volume of deci-normal dichromate solution would be reduce by 1 g of tin.
27. 5 g sample of brass was dissolved in one litre dil.  $\text{H}_2\text{SO}_4$ . 20 mL of this solution were mixed with KI, liberating  $\text{I}_2$  and  $\text{Cu}^+$  and the  $\text{I}_2$  required 20 mL of 0.0327 N hypo solution for complete titration. Calculate the percentage of Cu in the alloy.
28. One g of impure sodium carbonate is dissolved in water and the solution is made up to 250 mL. To 50 mL of this made up solution, 50 mL of 0.1 N – HCl is added and the mix after shaking well required 10 mL of 0.16 N – NaOH solution for complete titration. Calculate the % purity of the sample.
29. What amount of substance containing 60% NaCl, 37% KCl should be weighed out for analysis so that after the action of 25 mL of 0.1 N  $\text{AgNO}_3$  solution, excess of  $\text{Ag}^+$  is back titrates with 5 mL of  $\text{NH}_4\text{SCN}$  solution. Given that 1 mL of  $\text{NH}_4\text{SCN} = 1.1 \text{ mL of AgNO}_3$ .
30. A bottle labelled with "12 V  $\text{H}_2\text{O}_2$ " contain 700 mL solution. If a student mix 300 mL water in it what is the g/litre strength & normality and volume strength of final solution.
31. 0.84 g iron ore containing x percent of iron was taken in a solution containing all the iron in ferrous condition. The solution required x mL of a dichromatic solution for oxidizing the iron content to ferric state. Calculate the strength of dichromatic solution.
32. The neutralization of a solution of 1.2 g of a substance containing a mixture of  $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ ,  $\text{KHC}_2\text{O}_4 \cdot \text{H}_2\text{O}$  and different impurities of a neutral salt consumed 18.9 mL of 0.5 N NaOH solution. On titration with  $\text{KMnO}_4$  solution, 0.4 g, of the same substance needed 21.55 mL of 0.25 N  $\text{KMnO}_4$ . Calculate the % composition of the substance.
33. 50 g of a sample of  $\text{Ca(OH)}_2$  is dissolved in 50 mL of 0.5 N HCl solution. The excess of HCl was titrated with 0.3 N – NaOH. The volume of NaOH used was 20cc. Calculate % purity of  $\text{Ca(OH)}_2$ .
34. 50 mL of an aqueous solution of  $\text{H}_2\text{O}_2$  were treated with an excess of KI solution and dilute  $\text{H}_2\text{SO}_4$ , the liberated iodine required 20 mL of 0.1 N  $\text{Na}_2\text{S}_2\text{O}_3$  solution for complete interaction. Calculate the concentration of  $\text{H}_2\text{O}_2$  in g/l.
35. 100 kg hard water contains 5 g  $\text{MgSO}_4$ . Find hardness.
36. One litre hard water contains 1 mg  $\text{CaCl}_2$  and 1 mg  $\text{MgSO}_4$ . Find hardness.
37. Calculate the amount of lime  $\text{Ca(OH)}_2$  required to remove the hardness in 60 litre of pond water containing 1.62 mg of calcium bicarbonate per 100 mL of water.

38. 10 g sample of bleaching powder was dissolved into water to make the solution one litre. To this solution 35 mL of 1.0 M Mohr salt solution was added containing enough  $\text{H}_2\text{SO}_4$ . After the reaction was complete, the excess Mohr salt required 30 mL of 0.1 M  $\text{KMnO}_4$  for oxidation. Find out the % of available  $\text{Cl}_2$  approximately is (mol wt. 71).
39. Calculate the amount (in milligrams) of  $\text{SeO}_3^{-2}$  in solution on the basis of following data 20 mL of M/60 solution of  $\text{KBrO}_3$  was added to a definite volume of  $\text{SeO}_3^{-2}$  solution. The bromine evolved was removed by boiling and excess of  $\text{KBrO}_3$  was back titrated with 5 mL of M/25 solution of  $\text{NaAsO}_2$ . The reactions are given below. (Atomic mass of K = 39, Br = 80, As = 75, Na = 23, O = 16, Se = 79)
- (a)  $\text{SeO}_3^{-2} + \text{BrO}_3^- + \text{H}^+ \longrightarrow \text{SeO}_4^{-2} + \text{Br}_2 + \text{H}_2\text{O}$
- (b)  $\text{BrO}_3^- + \text{AsO}_2^- + \text{H}_2\text{O} \longrightarrow \text{Br}^- + \text{AsO}_4^{-3} + \text{H}^+$
40. Calculate the hardness of water sample which contains 0.001 mol  $\text{MgSO}_4$  per litre of water.
41. A solution of a 0.4 g sample of  $\text{H}_2\text{O}_2$  reacted with 0.632 g of  $\text{KMnO}_4$  in the presence of sulphuric acid. Calculate the percentage purity of the sample of  $\text{H}_2\text{O}_2$ .
42. 5 litre of a solution of  $\text{H}_2\text{O}_2$  with x N strength is diluted to 5.5 litre. This 5.5 litre  $\text{H}_2\text{O}_2$  solution gives 28 litre  $\text{O}_2$  at NTP. Find the value of x.
43. A 1.0 g sample of  $\text{Fe}_2\text{O}_3$  solid of 55.2% purity is dissolved in acid and reduced by heating the solution with zinc dust. The resultant solution is cooled and made upto 100.0 mL. An aliquot of 25.0 mL of this solution requires 17.0 mL of 0.0167 M solution of an oxidant for titration. Calculate the number of moles of electrons taken up by the oxidant in the reaction of the above titration.
44. 25 mL of a solution containing HCl was treated with excess of M/5  $\text{KIO}_3$  and KI solution of unknown concentration where  $\text{I}_2$  liberated is titrated against a standard solution of 0.021 M  $\text{Na}_2\text{S}_2\text{O}_3$  solution whose 24 mL were used up. Find the strength of HCl and volume of  $\text{KIO}_3$  solution consumed :
45. 0.6213 g of sample contains an unknown amount of  $\text{As}_2\text{O}_3$ . The sample was treated with HCl resulting information of  $\text{AsCl}_3$  (g) which was distilled into a beaker of water. The hydrolysis reaction is as follows :
- $$\text{AsCl}_3 + 2\text{H}_2\text{O} \rightarrow \text{HAsO}_2 + 3\text{H}^+ + 3\text{Cl}^-$$
- The amount of  $\text{HAsO}_2$  was determined by titration with 0.04134 M  $\text{I}_2$ , requiring 23.04 mL to reach the equivalence point. The redox products in the titration were  $\text{H}_3\text{AsO}_4$  and  $\text{I}^-$ . Find the amount of  $\text{KMnO}_4$  needed to oxidize As in  $\text{As}_2\text{O}_3$  to its maximum possible oxidation state in acidic medium.
46. A sample of steel weighing 0.6 g and containing S as an impurity was burnt in a stream of  $\text{O}_2$ , when S was converted to its oxide  $\text{SO}_2$ .  $\text{SO}_2$  was then oxidized to  $\text{SO}_4^{2-}$  by using  $\text{H}_2\text{O}_2$  solution containing 30 mL of 0.04 M NaOH. 22.48 mL of 0.024 M HCl was required to neutralize the base remaining after oxidation. Calculate the % of S in the sample :
47. 1.2475 g of crystalline copper sulphate was dissolved in water and excess of KI was added. The liberated iodine consumed 50 mL N/10  $\text{Na}_2\text{S}_2\text{O}_3$  solution to reach the end point of the titration. Calculate the number of water molecules of hydration in crystalline copper sulphate salt.
48. A 1g sample of  $\text{K}_2\text{Cr}_2\text{O}_7$  containing some inert material was entirely reduced with conc. HCl. The chlorine liberated was passed through hot solution of NaOH at  $80^\circ\text{C}$ , and it completely disproportionates to form  $\text{ClO}_3^-$  and  $\text{Cl}^-$ . This  $\text{NaClO}_3$  was isolated and its reduction with KI (aq) liberated iodine, giving  $\text{Cl}^-$ . The iodine thus liberated required 100 mL of decinormal hypo solution for complete titration. What is the percentage purity of the dichromate sample?

## REDOX REACTION AND EQUIVALENT CONCEPT

49. 2.5g of mixture of crystalline oxalic acid ( $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ ) and sodium oxalate ( $\text{Na}_2\text{C}_2\text{O}_4$ ) was dissolved in 100 mL of water. 50 mL of this solution was titrated against N/10 NaOH solution when 119.05 mL of the base was found necessary to reach the end point with phenolphthalein as the indicator. 1g of the mixture was dissolved in water and the solution titrated against N/10  $\text{KMnO}_4$  in the presence of dil.  $\text{H}_2\text{SO}_4$ . What is the volume of  $\text{KMnO}_4$  needed for getting the end point with 0.5g of the mixture?
50. In the presence of fluoride ion,  $\text{Mn}^{2+}$  can be titrated with  $\text{MnO}_4^-$ , both reactants being converted to a complex of Mn(III). A 0.545 g sample containing  $\text{Mn}_3\text{O}_4$  was dissolved and all manganese was converted to  $\text{Mn}^{2+}$ . Titration in the presence of fluoride ion consumed 31.1 mL of  $\text{KMnO}_4$  that was 0.177 N against oxalate.
- (a) write a balanced chemical equation for the reaction, assuming that the complex is  $\text{MnF}_4^-$ .
- (b) what was the % of  $\text{Mn}_3\text{O}_4$  in the sample ?
51. A mixture of two gases,  $\text{H}_2\text{S}$  and  $\text{SO}_2$  is passed through three beakers successively. The first beaker contains  $\text{Pb}^{2+}$  ions, which absorbs  $\text{S}^{2-}$  forming  $\text{PbS}$ . The second beaker contains 25 mL of 0.0396 N  $\text{I}_2$  to oxidize  $\text{SO}_2$  to  $\text{SO}_4^{2-}$ . The third contains 10 mL of 0.0345 N thiosulphate solution to retain any  $\text{I}_2$  carried over from the second absorber. A 25 L gas sample was passed through the apparatus followed by an additional amount of  $\text{N}_2$  to sweep last traces of  $\text{SO}_2$  from first and second absorber. The solution from the first absorber was made acidic and treated with 20 mL of 0.0066 M  $\text{K}_2\text{Cr}_2\text{O}_7$  which converted  $\text{S}^{2-}$  to  $\text{SO}_2$ . The excess dichromate was reacted with solid KI and the liberated iodine required 7.45 mL of 0.0345 N  $\text{Na}_2\text{S}_2\text{O}_3$  solution. The solutions in the second and third absorbers were combined and the resultant iodine was titrated with 2.44 mL of the same thiosulphate solution. Calculate the concentrations of  $\text{SO}_2$  and  $\text{H}_2\text{S}$  in mg/L of the sample :
52. 1 g of a moist sample of a mixture of  $\text{KClO}_3$  and  $\text{KCl}$  was dissolved in water and made upto 250 mL. 25 mL of this solution was treated with  $\text{SO}_2$  to reduced chlorate into chloride and the excess  $\text{SO}_2$  was boiled off. When the total chloride was precipitated, 0.1435 g of  $\text{AgCl}$  was obtained. In another experiment 25 mL of the original solution was treated with 30 mL of 0.2 N solution of  $\text{FeSO}_4$  and unreacted  $\text{FeSO}_4$  required 37.5 mL of 0.08 N solution of an oxidizing agent for complete oxidation. Calculate the molar ratio of chlorate and chloride in the given mixture.  $\text{Fe}^{2+}$  reacts with  $\text{ClO}_3^-$  according to equation :
- $$\text{ClO}_3^- + 6\text{Fe}^{2+} + 6\text{H}^+ \rightarrow \text{Cl}^- + 6\text{Fe}^{3+} + 3\text{H}_2\text{O}$$
- Also calculate the mass percent of moisture present in the moist sample.
53. A 1 g sample containing NaOH as the only basic substance and some inert impurity was left exposed to atmosphere for a very long time so that part of NaOH got converted into  $\text{Na}_2\text{CO}_3$  by absorbing  $\text{CO}_2$  from atmosphere. The resulting sample was dissolved in water and volume made upto 100 mL. A 20 mL portion of this solution required 16 mL 0.25 M HCl solution to reach the equivalence point when methyl orange was used as indicator. In a separate analysis, 20 mL portion of the same solution was taken along with phenolphthalein indicator and mixed with 50 mL of 0.1 M HCl solution. An additional 9.00 mL 0.1 M  $\text{Ba}(\text{OH})_2$  solution was required to just restore the pink colour of solution. Determine mass percentage of NaOH in the original sample and mass percentage of  $\text{Na}_2\text{CO}_3$  in the sample after exposure to atmosphere.

54. A steel sample is to be analysed for Cr and Mn simultaneously. By suitable treatment the Cr is oxidized to  $\text{Cr}_2\text{O}_7^{2-}$  and the Mn to  $\text{MnO}_4^-$ . A 10.00 g sample of steel is used to produce 250.0 mL of a solution containing  $\text{Cr}_2\text{O}_7^{2-}$  and  $\text{MnO}_4^-$ . A 10.00 mL portion of this solution is added to a  $\text{BaCl}_2$  solution and by proper adjustment of the acidity, the chromium is completely precipitated as  $\text{BaCrO}_4$ ; 0.0549 g is obtained. A second 10.00 mL portion of this solution requires exactly 15.95 mL of 0.0750 M standard  $\text{Fe}^{2+}$  solution for its titration (in acid solution). Calculate the % of Mn and % of Cr in the steel sample.
55. 1.16 g  $\text{CH}_3(\text{CH}_2)_n\text{COOH}$  was burnt in excess air and the resultant gases ( $\text{CO}_2$  and  $\text{H}_2\text{O}$ ) were passed through excess NaOH solution. The resulting solution was divided in two equal parts. One part required 50 mL of 1 N HCl for neutralization using phenolphthalein as indicator. Another part required 80 mL of 1 N HCl for neutralization using methyl orange as indicator. Find the value of n and the amount of excess NaOH solution taken initially.
56. A 1.5 g sample containing oxalic acid and some inert impurity was dissolved in enough water and volume made up to 250 mL. A 20 mL portion of this solution was then mixed with 30 mL of an alkali solution. The resulting solution was then treated with stoichiometric amount of  $\text{CaCl}_2$  just needed for precipitation of oxalate as  $\text{CaC}_2\text{O}_4$ . Solution was filtered off and filtrate was finally titrated against 0.1 M HCl solution. 8.0 mL of acid was required to reach the equivalence point. At last, the above neutral solution was treated with excess of  $\text{AgNO}_3$  solution and AgCl obtained was washed, dried and weighed to be 0.4305 g. Determine mass percentage of oxalic acid in the original sample :



## Exercise # 5

## Part # I

## [Previous Year Questions] [AIEEE/JEE-MAIN]

- The oxidation state of Cr in  $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]^+$  is - [AIEEE-05]  
 (1) +2 (2) +3 (3) 0 (4) +1
- The oxidation state of chromium in the final product formed by the reaction between KI and acidified potassium dichromate solution is - [AIEEE-05]  
 (1) +6 (2) +4 (3) +3 (4) +2
- Which of the following chemical reaction depicts the oxidizing behaviour of  $\text{H}_2\text{SO}_4$  ? [AIEEE-06]  
 (1)  $\text{Ca}(\text{OH})_2 + \text{H}_2\text{SO}_4 \rightarrow \text{CaSO}_4 + 2\text{H}_2\text{O}$  (2)  $\text{NaCl} + \text{H}_2\text{SO}_4 \rightarrow \text{NaHSO}_4 + \text{HCl}$   
 (3)  $2\text{PCl}_5 + \text{H}_2\text{SO}_4 \rightarrow 2\text{POCl}_3 + 2\text{HCl} + \text{SO}_2\text{Cl}_2$  (4)  $2\text{HI} + \text{H}_2\text{SO}_4 \rightarrow \text{I}_2 + \text{SO}_2 + 2\text{H}_2\text{O}$
- Which of the following reactions is an example of a redox reaction? [JEE Main-2017]  
 (1)  $\text{XeF}_4 + \text{O}_2\text{F}_2 \rightarrow \text{XeF}_6 + \text{O}_2$  (2)  $\text{XeF}_2 + \text{PF}_5 \rightarrow [\text{XeF}]^+ + \text{PF}_6^-$   
 (3)  $\text{XeF}_6 + \text{H}_2\text{O} \rightarrow \text{XeOF}_4 + 2\text{HF}$  (4)  $\text{XeF}_6 + 2\text{H}_2\text{O} \rightarrow \text{XeO}_2\text{F}_2 + 4\text{HF}$
- Hydrogen peroxide oxidises  $[\text{Fe}(\text{CN})_6]^{4-}$  to  $[\text{Fe}(\text{CN})_6]^{3-}$  in acidic medium but reduces  $[\text{Fe}(\text{CN})_6]^{3-}$  to  $[\text{Fe}(\text{CN})_6]^{4-}$  in alkaline medium. The other products formed are, respectively: [JEE Main-2018]  
 (1)  $(\text{H}_2\text{O} + \text{O}_2)$  and  $(\text{H}_2\text{O} + \text{OH}^-)$  (2)  $\text{H}_2\text{O}$  and  $(\text{H}_2\text{O} + \text{O}_2)$   
 (3)  $\text{H}_2\text{O}$  and  $(\text{H}_2\text{O} + \text{OH}^-)$  (4)  $(\text{H}_2\text{O} + \text{O}_2)$  and  $\text{H}_2\text{O}$

## Part # II

## [Previous Year Questions] [IIT-JEE ADVANCED]

- In basic medium  $\text{I}^-$  oxidises by  $\text{MnO}_4^-$ . In this process  $\text{I}^-$  replaces by : [JEE 2004]  
 (A)  $\text{IO}_3^-$  (B)  $\text{I}_2$  (C)  $\text{IO}_4^-$  (D)  $\text{IO}^-$
- Amongst the following, the pair having both the metals in their highest oxidation state is : [JEE 2004]  
 (A)  $[\text{Fe}(\text{CN})_6]^{3-}$  and  $[\text{Co}(\text{CN})_6]^{3-}$  (B)  $[\text{CrO}_2\text{Cl}_2]$  and  $[\text{MnO}_4^-]$   
 (C)  $\text{TiO}_2$  and  $\text{MnO}_2$  (D)  $[\text{MnCl}_4]^{2-}$  and  $[\text{NiF}_6]^{2-}$
- $\text{O}_3$  does not oxidise : [JEE 2005]  
 (A) KI (B)  $\text{FeSO}_4$  (C)  $\text{KMnO}_4$  (D)  $\text{K}_2\text{MnO}_4$
- A  $5.0 \text{ cm}^3$  solution of  $\text{H}_2\text{O}_2$  liberates 0.508 g of iodine from an acidified KI solution. Calculate the strength of  $\text{H}_2\text{O}_2$  solution in terms of volume strength at STP. [JEE' 1995]
- A 3.00 g sample containing  $\text{Fe}_3\text{O}_4$ ,  $\text{Fe}_2\text{O}_3$  and an inert impure substance, is treated with excess of KI solution in presence of dilute  $\text{H}_2\text{SO}_4$ . The entire iron is converted into  $\text{Fe}^{2+}$  along with the liberation of iodine. The resulting solution is diluted to 100 mL. A 20 mL of the diluted solution require 11 mL of 0.5 M  $\text{Na}_2\text{S}_2\text{O}_3$  solution to reduce the iodine present. A 50 mL of diluted solution after complete extraction of the iodine requires 12.80 mL of 0.25  $\text{KMnO}_4$  solution in dilute  $\text{H}_2\text{SO}_4$  medium for the oxidation of  $\text{Fe}^{2+}$ . Calculate the percentages of  $\text{Fe}_2\text{O}_3$  and  $\text{Fe}_3\text{O}_4$  in the original sample. [JEE 2000]
- One litre of a mixture of  $\text{O}_2$  and  $\text{O}_3$  at NTP was allowed to react with an excess of acidified solution of KI. The Iodine liberated required 40 mL of M/10 sodium thiosulphate solution for titration. What is the percent of ozone in the mixture ? Ultraviolet radiation of wavelength 300 nm can decompose ozone. Assuming that one photon can decompose one ozone molecule, how many photons would have been required for the complete decomposition of ozone in the original mixture? [JEE 97,5]

7. A sample of hard water contains 96 ppm of  $\text{SO}_4^{2-}$  and 183 ppm of  $\text{HCO}_3^-$ , with  $\text{Ca}^{2+}$  as the only cation. How many moles of CaO will be required to remove  $\text{HCO}_3^-$  from 1000 kg of this water? If 1000 kg of this water is treated with the amount of CaO calculate above, what will be the concentration (in ppm) of residual  $\text{Ca}^{2+}$  ions (Assume  $\text{CaCO}_3$  to be completely insoluble in water)? If the  $\text{Ca}^{2+}$  ions in one litre of the treated water are completely exchanged with hydrogen ions, what will be its pH (one ppm means one part of the substance in one million part of water, weight / weights)? [JEE' 1997]
8. An aqueous solution containing 0.10 g  $\text{KIO}_3$  (formula wt. 214.0) was treated with an excess of KI solution. The solution was acidified with HCl. The liberated  $\text{I}_2$  consumed 45.0 mL of thiosulphate solution to decolourise the blue starch – iodine complex. Calculate the molarity of the sodium thiosulphate solution. [JEE 1998]
9. How many millilitre of 0.5 M  $\text{H}_2\text{SO}_4$  are needed to dissolve 0.5 g of copper II carbonate? [JEE 1999]
10. Hydrogen peroxide solution (20 mL) reacts quantitatively with a solution of  $\text{KMnO}_4$  (20 mL) acidified with dilute  $\text{H}_2\text{SO}_4$ . The same volume of  $\text{KMnO}_4$  solution is just decolorized by 10 mL of  $\text{MnSO}_4$  in neutral medium simultaneously forming a dark brown precipitate of hydrated  $\text{MnO}_2$ . The brown precipitate is dissolved in 10 mL of 0.2 M sodium oxalate under boiling condition in the presence of dilute  $\text{H}_2\text{SO}_4$ . Write the balanced equations involved in the reactions and calculate the molarity of  $\text{H}_2\text{O}_2$ . [JEE 2001]
11. For the reaction :  

$$\text{I}^- + \text{ClO}_3^- + \text{H}_2\text{SO}_4 \longrightarrow \text{Cl}^- + \text{HSO}_4^- + \text{I}_2$$
 The correct statement(s) in the balanced equation is/are :  
 (A) Stoichiometric coefficient of  $\text{HSO}_4^-$  is 6 (B) Iodide is oxidized.  
 (C) Sulphur is reduced (D)  $\text{H}_2\text{O}$  is one of the products
12. Hydrogen peroxide in its reaction with  $\text{KIO}_4$  and  $\text{NH}_2\text{OH}$  respectively, is acting as a  
 (A) reducing agent, oxidising agent (B) reducing agent, reducing agent  
 (C) oxidising agent, oxidising agent. (D) oxidising agent, reducing agent
13. In dilute aqueous  $\text{H}_2\text{SO}_4$ , the complex diaquodioxalatoferrate(II) is oxidized by  $\text{MnO}_4^-$ . For this reaction, the ratio of the rate of change of  $[\text{H}^+]$  to the rate of change of  $[\text{MnO}_4^-]$  is. [JEE 2015]

MOCK TEST

SECTION - I : STRAIGHT OBJECTIVE TYPE

- The number of moles of  $\text{KMnO}_4$  that will be needed to react completely with one mole of ferrous oxalate in acid solution is  
(A)  $3/5$  (B)  $2/5$  (C)  $4/5$  (D) 1
- The number of moles of  $\text{KMnO}_4$  that will be needed to react with one mole of sulphite ions in acidic solution is  
(A)  $2/5$  (B)  $3/5$  (C)  $4/5$  (D) 1
- The normality of 0.3 M phosphorus acid ( $\text{H}_3\text{PO}_3$ ) is  
(A) 0.1 (B) 0.9 (C) 0.3 (D) 0.6
- An aqueous solution of 6.3 gm of oxalic acid dihydrate is made upto 250 ml. The volume of 0.1 N NaOH required to completely neutralize 10 ml of this solution is  
(A) 40 ml (B) 20 ml (C) 10 ml (D) 4 ml
- Consider a titration of potassium dichromate solution with acidified Mohr's salt solution using diphenylamine as indicator. The number of moles of Mohr's salt required per mole of dichromate is  
(A) 3 (B) 4 (C) 5 (D) 6

SECTION - II : MULTIPLE CORRECT ANSWER TYPE

- Consider the redox reaction  $2\text{S}_2\text{O}_3^{2-} + \text{I}_2 \longrightarrow \text{S}_4\text{O}_6^{2-} + 2\text{I}^-$ :  
(A)  $\text{S}_2\text{O}_3^{2-}$  gets reduced to  $\text{S}_4\text{O}_6^{2-}$  (B)  $\text{S}_2\text{O}_3^{2-}$  gets oxidised to  $\text{S}_4\text{O}_6^{2-}$   
(C)  $\text{I}_2$  gets reduced to  $\text{I}^-$  (D)  $\text{I}_2$  gets oxidised to  $\text{I}^-$
- There are two sample of HCl having molarity 1N and 0.25 N. Find volume of these sample taken in order to prepare 0.75 N HCl solution. (Assume no water is used) :  
(A) 20 mL, 10 mL (B) 100 mL, 50 mL (C) 40 mL, 20 mL (D) 50 mL, 25 mL
- To a 25 ml  $\text{H}_2\text{O}_2$  solution excess acidified solution of KI was added. The iodine liberated 20 ml of 0.3 N sodium thiosulphate solution. Use these data to choose the correct statements from the following :  
(A) The weight of  $\text{H}_2\text{O}_2$  present in 25 ml solution is 0.102 g  
(B) The molarity of  $\text{H}_2\text{O}_2$  solution is 0.12 M  
(C) The weight of  $\text{H}_2\text{O}_2$  present in 1 L of the solution is 0.816 g  
(D) The volume strength of  $\text{H}_2\text{O}_2$  is 1.344 L

SECTION - III : ASSERTION AND REASON TYPE

Each question has 5 choices (A), (B), (C), (D) and (E) out of which only one is correct.

- Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.  
(B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1.  
(C) Statement-1 is True, Statement-2 is False.  
(D) Statement-1 is False, Statement-2 is True.  
(E) Statement-1 and Statement-2 both are False.
- Statement-1 :** In the redox reaction  $8\text{H}^+(\text{aq}) + 4\text{NO}_3^- + 6\text{Cl}^- + \text{Sn}(\text{s}) \longrightarrow \text{SnCl}_6^{2-} + 4\text{NO}_2 + 4\text{H}_2\text{O}$ , the reducing agent is Sn(s).  
**Statement-2 :** In balancing half reaction,  $\text{S}_2\text{O}_3^{2-} \longrightarrow \text{S}(\text{s})$ , the number of electrons added on the left is 4.

## CHEMISTRY FOR JEE MAIN & ADVANCED


10. **Statement-1** : Among  $\text{Br}^-$ ,  $\text{O}_2^{2-}$ ,  $\text{H}^-$  and  $\text{NO}_3^-$ , the ions that cannot act as oxidising agents are  $\text{Br}^-$  and  $\text{H}^-$   
**Statement-2** :  $\text{Br}^-$  and  $\text{H}^-$  cannot be reduced.

### SECTION - IV : COMPREHENSION TYPE

Read the following comprehensions carefully and answer the questions.

#### Comprehension # 1

Chromium exists as  $\text{FeCr}_2\text{O}_4$  in the nature and it contains  $\text{Fe}_{0.95}\text{O}$  & other impurity. To obtain pure chromium from  $\text{FeCr}_2\text{O}_4$ , the ore is fused with  $\text{KOH}$  and oxygen is passed through the mixture when  $\text{K}_2\text{CrO}_4$  and  $\text{Fe}_2\text{O}_3$  are produced.



2 g of ore required 270 mL of  $\text{O}_2$  at 273 K and 1 atm for complete oxidation of ore.  $\text{K}_2\text{CrO}_4$  is precipitated as  $\text{BaCrO}_4$  when Barium salt is added. To remaining solution 10 mL of 1 M  $\text{K}_4\text{Fe}(\text{CN})_6$  is added when  $\text{Fe}^{3+}$  ions react with it to form  $\text{KFe}[\text{Fe}(\text{CN})_6]$ , often called 'Prussian Blue'. To determine excess of  $\text{K}_4\text{Fe}(\text{CN})_6$  in solution 7 mL of 0.2 N of  $\text{Fe}^{2+}$  is added when all the  $\text{K}_4\text{Fe}(\text{CN})_6$  is precipitated as  $\text{K}_2\text{Fe}[\text{Fe}(\text{CN})_6]$ .

11. Weight of  $\text{BaCrO}_4$  precipitated  
(A) 1.64 (B) 6.29 (C) 0.82 (D) 3.29
12. % by mass of  $\text{Fe}_{0.95}\text{O}$  in the ore  
(A) 9.6% (B) 10.1% (C) 8.55% (D) 20.2%
13. n factor for  $\text{Fe}_{0.95}\text{O}$   
(A) 0.9 (B) 0.85 (C)  $\frac{2}{0.95}$  (D) 1.8
14. Weight of impurities present in the ore  
(A) 0.421 (B) 0.123 (C) 0.341 (D) 0.206

### SECTION - V : MATRIX - MATCH TYPE

15. **Column I** **Column II**
- |   |  |
|---|--|
| (A) $\text{Sn}^{+2}$ + $\text{MnO}_4^-$ (acidic)<br>3.5 mole 1.2 mole                 | (p) Amount of oxidant available decides the number of electrons transfer                                   |
| (B) $\text{H}_2\text{C}_2\text{O}_4$ + $\text{MnO}_4^-$ (acidic)<br>8.4 mole 3.6 mole | (q) Amount of reductant available decides the number of electrons transfer                                 |
| (C) $\text{S}_2\text{O}_3^{2-}$ + $\text{I}_2$<br>7.2 mole 3.6 mole                   | (r) Number of electrons involved per mole of oxidant > Number of electrons involved per mole of reductant  |
| (D) $\text{Fe}^{+2}$ + $\text{Cr}_2\text{O}_7^{2-}$ (acidic)<br>9.2 mole 1.6 mole     | (s) Number of electrons involved per mole of oxidant < Number of electrons involved per mole of reductant. |

### SECTION - VI : SUBJECTIVE TYPE

16.  $\text{H}_2\text{O}_2$  is reduced rapidly by  $\text{Sn}^{2+}$ , the products being  $\text{Sn}^{4+}$  & water.  $\text{H}_2\text{O}_2$  decomposes slowly at room temperature to yield  $\text{O}_2$  & water. Calculate the volume of  $\text{O}_2$  produced at 273K & 1.00 atm when 200g of 10.0 % by mass  $\text{H}_2\text{O}_2$  in water is treated with 88.2 ml of 1 M  $\text{Sn}^{2+}$  & then the mixture is allowed to stand until no further reaction occurs.
17. 80 gm of a sample of Anhydrous  $\text{CuSO}_4$  was dissolved in water and made to 250ml. 25 ml of this solution after taking usual precautions was treated with a little excess of  $\text{KI}$  solution. A white ppt. of  $\text{Cu}_2\text{I}_2$  and iodine was evolved. The iodine so evolved required 40 ml of 1 M of hypo solution. What is the approximate purity of  $\text{CuSO}_4$  solution. [Mol. wt. of  $\text{CuSO}_4 = 160$ ]

ANSWER KEY

EXERCISE - 1

1. D 2. A 3. C 4. C 5. B 6. D 7. D 8. C 9. A 10. B 11. C 12. A 13. C  
 14. C 15. A 16. B 17. A 18. (i) A (ii) A (iii) A (iv) D 19. A 20. D 21. B 22. B  
 23. A 24. D 25. B 26. A 27. A 28. B 29. B 30. C 31. D 32. B 33. B 34. C 35. A  
 36. A 37. D 38. D 39. D 40. C 41. D 42. A 43. D 44. A 45. B 46. B 47. A 48. A  
 49. B 50. B

EXERCISE - 2 : PART # I

1. A,C 2. A,B,C,D 3. A,C,D 4. A,B,D 5. B,C 6. A,B,C,D  
 7. A,B,D 8. B,D 9. B,C 10. A,B,D 11. A,B,D 12. A,C,D  
 13. B,C,D 14. A,B,C 15. B,C 16. D 17. C 18. A,B  
 19. A,B 20. A,B,D

PART # II

- 1 B 2 A 3 A 4 A 5 B 6 A 7 B 8 A 9 A 10 A 11 B 12 D 13 C

EXERCISE - 3 : PART # I

1.  $A \rightarrow (q), B \rightarrow (p), C \rightarrow (s), D \rightarrow (r)$  2.  $A \rightarrow (s), B \rightarrow (p), C \rightarrow (q), D \rightarrow (r)$   
 3.  $A \rightarrow (s), B \rightarrow (q), C \rightarrow (r), D \rightarrow (p)$  4.  $A \rightarrow (p), B \rightarrow (r,s), C \rightarrow (r), D \rightarrow (r, q)$   
 5.  $A \rightarrow (p, s), B \rightarrow (s), C \rightarrow (p, q), D \rightarrow (r)$  6.  $A \rightarrow (p, s), B \rightarrow (q, r), C \rightarrow (p, q, s), D \rightarrow (r)$   
 7.  $A \rightarrow (p, r), B \rightarrow (q, r), C \rightarrow (p, q, r), D \rightarrow (q, r)$

PART # II

- Comprehension #1: 1. A 2. B 3. B 4. B  
 Comprehension #2: 1. B 2. C 3. C 4. B  
 Comprehension #3: 1. C 2. A 3. C 4. B  
 Comprehension #4: 1. C 2. D 3. B  
 Comprehension #5: 1. C 2. B 3. C

EXERCISE - 5 : PART # I

1. 2      2. 3      3. 4      4. 1    5. 2

PART # II

1. A    2. B    3. C    4. 4.48    5.  $\text{Fe}_2\text{O}_3 = 49.33\%$ ,  $\text{Fe}_3\text{O}_4 = 34.8\%$     6. 6.57%  $\text{O}_3$  (by weight),  $1.2 \times 10^{21}$  photons  
7. 1.5, 40 ppm,  $\text{pH} = 2.6989$     8. 0.0623 M    9. 8.097 mL    10. 0.1 M    11. (A), (B), (D)    12. (A)    13. 8

MOCK TEST

1. A    2. A    3. D    4. A    5. D    6. B,C    7. A,B,C,D    8. A,B,D    9. C    10. B    11. D    12. B  
13. B    14. C    15.  $\text{A} \rightarrow (\text{p}, \text{r}), \text{B} \rightarrow (\text{q}, \text{r}), \text{C} \rightarrow (\text{p}, \text{q}, \text{r}), \text{D} \rightarrow (\text{q}, \text{r})$     16. 5.6L    17. 80%