

Chapter 11

Dual Nature of Radiation and Matter

EXERCISE

Question 1:

Find the

- maximum frequency, and
- Minimum wavelength of X-rays produced by 30 kV electrons.

Solution 1:

a) We know,

$$K_{\max} = h\nu - \Phi_0$$

$$\text{Or } eV_0 = h\nu - \Phi_0$$

$$\nu_{\max} = eV_0 / h$$

Substituting the required values

$$\nu_{\max} = (1.6 \times 10^{-19}) \cdot (30000) / (6.63 \times 10^{-34})$$

$$= 7.24 \times 10^{18} \text{ Hz}$$

$$\begin{aligned} \text{(b) By the relation } \lambda &= c / \nu_{\max} \\ &= 0.041 \text{ nm} \end{aligned}$$

Question 2:

The work function of caesium metal is 2.14 eV, When light of frequency 6×10^{14} Hz is incident on the metal surface, photoemission of electrons occurs.

- What is the maximum kinetic energy of the emitted electrons,
- Stopping potential, and
- Maximum speed of the emitted photoelectrons?

Solution 2:

Given:

$$\Phi_0 = 2.14 \text{ eV}$$

$$\nu = 6 \times 10^{14} \text{ Hz}$$

a) Maximum kinetic energy is the difference between the photon energy and the work function of the metal. For caesium it is given to be 2.14 eV

Using the expression

$$K_{\max} = h\nu - \Phi_0$$

Substitution yields

Chapter 11

Dual Nature of Radiation and Matter

$$K_{\max} = \left[(6.63 \times 10^{-34}) \cdot (6 \times 10^{14}) \right] - \left[(2.114) \cdot (1.6 \times 10^{19}) \right]$$

$$= 5.54 \times 10^{-20} \text{ J}$$

$$= 0.346 \text{ eV}$$

b) Stopping potential can be deduced by maximum kinetic energy, by the expression

$$eV_0 = K_{\max}$$

$$\text{Or } V_0 = K_{\max} / e$$

Putting the required values

$$V_0 = 5.54 \times 10^{-20} / 1.6 \times 10^{-19}$$

$$= 0.34 \text{ V}$$

c) Maximum speed of photoelectron is given by the maximum kinetic energy it possesses,

$$K_{\max} = 5.54 \times 10^{-20}$$

$$\text{Or } mv^2 / 2 = 5.54 \times 10^{-20}$$

$$\text{Or } v_{\max} = 344 \text{ km / s}$$

Question 3:

The photoelectric cut-off voltage in a certain experiment is 1.5 V. What is the maximum kinetic energy of photoelectrons emitted?

Solution 3:

Maximum kinetic energy,

$$K_{\max} = eV_0$$

Putting values and solving

$$= 2.4 \times 10^{-19} \text{ J}$$

Question 4:

Monochromatic light of wavelength 632.8 nm is produced by a helium-neon laser. The power emitted is 9.42 m W.

a) Find the energy and momentum of each photon in the light beam,

b) How many photons per second, on the average, arrive at a target irradiated by this beam? (Assume the beam to have uniform cross-section which is less than the target area), and

c) How fast does a hydrogen atom have to travel in order to have the same momentum as that of the photon?

Solution 4:

Given:

Wavelength of monochromatic light

$$\lambda = 632.8 \text{ nm}$$

Power of He-be laser

$$P = 9.42 \text{ mW}$$

(a) Energy of a photon is given by

$$E = h\nu$$

$$\text{Or } E = hc / \lambda$$

Which gives

$$E = 3.14 \times 10^{-19} \text{ J}$$

Now momentum of a photon

$$p = h / \lambda$$

$$p = 1.05 \times 10^{-27} \text{ kg m / s}$$

b) For a beam of uniform cross-section having cross-sectional area less than target area

$$P = E \times N$$

where

P = power emitted

E = energy of photon

N = number of photons

Therefore

$$N = P / E$$

Substitution gives

$$N = 3 \times 10^{16} \text{ photons / second}$$

(c) Momentum of He-Ne laser = $1.05 \times 10^{-27} \text{ kg m / s}$

For this much momentum of a hydrogen atom

$$mv = 1.05 \times 10^{-27}$$

$$\text{Or } v = 1.05 \times 10^{-27} / 1.6 \times 10^{-27}$$

$$\text{Or } v = 0.63 \text{ m / s}$$

The required speed for hydrogen atom is 0.63 m / s.

Question 5:

The energy flux of sunlight reaching the surface of the earth is $1.388 \times 10^3 \text{ W/m}^2$. How many photons (nearly) per square meter are incident on the Earth per second? Assume that the photons in the sunlight have an average wavelength of 550 nm.

Solution 5:

Given:

Energy flux = 1388 W / sq. m

wavelength = 550 nm

Energy of photon = hc / λ

$$= 3.61 \times 10^{-19} \text{ J}$$

So no. of photons = P / E

$$= 4 \times 10^{21} \text{ Photons/ } m^2 s$$

Question 6:

In an experiment on photoelectric effect, the slope of the cut-off voltage versus frequency of incident light is found to be 4.12×10^{-15} V s. Calculate the value of Planck's constant.

Solution 6:

$$\text{Slope} = 4.12 \times 10^{-15}$$

$$\text{Or } h/e = 4.12 \times 10^{-15}$$

Putting value of e and solving

$$h = 6.592 \times 10^{-34} \text{ JS}$$

Question 7:

A 100W sodium lamp radiates energy uniformly in all directions. The lamp is located at the centre of a large sphere that absorbs all the sodium light which is incident on it. The wavelength of the sodium light is 589 nm. (a) What is the energy per photon associated with the sodium light? (b) At what rate are the photons delivered to the sphere?

Solution 7:

Given:

$$P = 100 \text{ W}$$

$$\lambda = 589 \text{ nm}$$

Energy per photon = hc / λ

$$= 3.37 \times 10^{-19} \text{ J}$$

b) No. of photons per = P/E

$$= 3 \times 10^{20} \text{ Photons/ second}$$

Question 8:

The threshold frequency for a certain metal is 3.3×10^{14} Hz. If light of frequency 8.2×10^{14} Hz is incident on the metal, predict the cutoff-voltage for the photoelectric emission.

Solution 8:

Given:

$$\nu_0 = 3.3 \times 10^{14} \text{ Hz}$$

$$\nu = 8.2 \times 10^{14} \text{ Hz}$$

Cut-off voltage

$$eV_0 = h(\nu - \nu_0)$$

Substituting the values

$$V_0 = 2.03 \text{ eV}$$

Question 9:

The work function for a certain metal is 4.2 eV. Will this metal give photoelectric emission for incident radiation of wavelength 330nm?

Solution 9:

$$\phi_0 = 4.2 \text{ eV}$$

$$\text{Or } \nu_0 h = 4.2$$

$$\text{Or } \nu_0 = 1.01 \times 10^{15} \text{ Hz}$$

Incident wavelength = 330nm

$$\text{Or } \nu = 6 \times 10^{14} \text{ Hz}$$

Since ν is less than ν_0 , there will be no photoelectric emission.

Question 10:

Light of frequency 7.21×10^{14} Hz is incident on a metal surface. Electrons with a maximum speed of 6.0×10^5 m/s are ejected from the surface. What is the threshold frequency for photoemission of electrons?

Solution 10:

Given:

Chapter 11

Dual Nature of Radiation and Matter

$$v = 7.21 \times 10^{14} \text{ Hz}$$

$$v = 6 \times 10^5 \text{ m/s}$$

$$\text{Kinetic energy} = h(v - v_0)$$

$$\text{Or } mv^2 / 2 = h(v - v_0)$$

Solving the above equation

$$v_0 = 4.73 \times 10^{14} \text{ Hz}$$

Question 11:

Light of wavelength 488 nm is produced by an argon laser which is used in the photoelectric effect. When light from this spectral line is incident on the emitter, the stopping (cut-off) potential of photoelectrons is 0.38 V. Find the work function of the material from which the emitter is made.

Solution 11:

Given:

$$\lambda = 488 \text{ nm}$$

$$V_0 = 0.38 \text{ V}$$

We Know

$$eV_0 = hv - \Phi_0$$

Or

$$\Phi_0 = hc / \lambda - eV_0$$

$$= 3.46 \times 10^{-19} \text{ J}$$

$$= 2.16 \text{ eV}$$

Question 12:

Calculate the

- a) momentum, and
- b) de Broglie wavelength of the electrons accelerated through a potential difference of 56 V.

Solution 12:

a) Using the relation

$$p = \sqrt{2meV}$$

$$= 4.04 \times 10^{-24} \text{ kg m/s}$$

b) And,

$$\lambda = h / p$$

$$= 0.164 \text{ nm}$$

Question 13:

What is the

a) momentum,

b) speed, and

c) de Broglie wavelength of an electron with kinetic energy of 120 eV.

Solution 13:

a) $p^2 = 2mK$

Or $p = 5.91 \times 10^{-24} \text{ kg m/s}$

b) Speed

$$mv^2 / 2 = K$$

$$mv^2 / 2 = 1.92 \times 10^{-17}$$

$$v = 6.5 \times 10^6 \text{ m/s}$$

c) $\lambda = h / p = 0.112 \text{ nm}$

Question 14:

The wavelength of light from the spectral emission line of sodium is 589 nm. Find the kinetic energy at which

a) an electron, and

b) a neutron, would have the same de Broglie wavelength.

Solution 14:

Given:

$$\lambda = 589 \text{ nm}$$

For this λ ,

$$p = h / \lambda$$

$$= 1.12 \times 10^{-17} \text{ kg m/s}$$

(a) For electron,

$$K = p^2 / 2m_e$$

$$= 6.9 \times 10^{-25} \text{ J}$$

(b) For neutron,

$$K = p^2 / 2m_n$$

$$= 3.7 \times 10^{-28} \text{ J}$$

Question 15:

What is the de Broglie wavelength of

- a) a bullet of mass 0.040 kg travelling at the speed of 1.0 km/s,
- b) a ball of mass 0.060 kg moving at a speed of 1.0 m/s, and
- c) a dust particle of mass 1.0×10^{-9} kg drifting with a speed of 2.2 m/s?

Solution 15:

a) $m = 0.04 \text{ kg}$

$$v = 1 \text{ km / s}$$

$$\lambda = h / mv$$

$$= 1.66 \times 10^{-35} \text{ m}$$

b) $m = 0.06 \text{ kg}$

$$v = 1 \text{ m / s}$$

$$\lambda = 1.1 \times 10^{-32} \text{ m}$$

c) $m = 1 \times 10^{-9} \text{ kg}$

$$v = 2.2 \text{ m / s}$$

$$\lambda = 3 \times 10^{-25} \text{ m}$$

Question 16:

An electron and a photon each have a wavelength of 1.00 nm. Find

- a) Their momenta,
- b) the energy of the photon, and
- c) the kinetic energy of electron.

Solution 16:

a) Momentum of electron

$$p = h / \lambda$$

$$= 6.63 \times 10^{-25} \text{ kg m / s}$$

Momentum of photon

$$p = h / \lambda$$

$$= 6.63 \times 10^{-25} \text{ kg m / s}$$

b) Energy of photon

$$E = mc^2 \dots (1)$$

Now,

$$\lambda = h / mc$$

$$\text{Or } m = h / c\lambda$$

Substituting value of m in eq. (1)

$$E = hc / \lambda$$

$$= 1.24 \text{ keV}$$

c) Kinetic energy of electron

$$K = p^2 / 2m$$

$$= 2.41 \times 10^{-19} \text{ J}$$

$$= 1.51 \text{ eV}$$

Question 17:

For what kinetic energy of a neutron will the associated de Broglie wavelength be $1.40 \times 10^{-10} \text{ m}$?

b) Also find the de Broglie wavelength of a neutron, in thermal equilibrium with matter, having an average kinetic energy of $(3/2) k T$ at 300 K.

Solution 17:

Given:

$$\lambda = 1.4 \times 10^{-10} \text{ m}$$

a) Now

$$K = p^2 / 2m$$

$$\text{And } p = h / \lambda$$

Therefore

$$K = h^2 / 2m\lambda^2$$

$$\text{Or } K = 6.686 \times 10^{-21} \text{ J}$$

b) Kinetic Energy

$$K = (3/2)kT$$

Where

$$k = \text{Boltzmann constant} = 1.381 \times 10^{-23} \text{ J / K}$$

$$T = 300\text{K}$$

So

$$K = 6.21 \times 10^{-21} \text{ J}$$

Now

$$\lambda = \frac{h}{\sqrt{2mK}} = 0.145 \text{ nm}$$

Question 18:

Show that the wavelength of electromagnetic radiation is equal to the de Broglie wavelength of its quantum (photon).

Solution 18:

For a photon,

$$p = hv / c$$

Therefore,

$$h / p = c / v = \lambda$$

That is, the de Broglie wavelength of a photon equals the wavelength of electromagnetic radiation of which the photon is a quantum of energy and momentum.

Question 19:

What is the de Broglie wavelength of a nitrogen molecule in air at 300K? Assume that the molecule is moving with the root-mean square speed of molecules at this temperature. (Atomic mass of nitrogen = 14.0076 u)

Solution 19:

Given:

$$T = 300 \text{ K}$$

$$m = 14.0076 \times 1.661 \times 10^{-27} \text{ kg}$$

$v = rms$ speed of molecules at T

$$K = 1.5kT$$

Substitution results

$$K = 6.21 \times 10^{-21} \text{ J}$$

$$\text{Now } \lambda = \frac{h}{\sqrt{2mK}}$$

Which on solution gives

$$\lambda = 0.038 \text{ nm}$$

Additional Exercise

Question 20:

- (a) Estimate the speed with which electrons emitted from a heated emitter of an evacuated tube impinge on the collector maintained at a potential difference of 500 V with respect to the emitter. Ignore the small initial speeds of the electrons. The specific charge of the electron, i.e., its e/m is given to be $1.76 \times 10^{11} \text{ C kg}^{-1}$.
- (b) Use the same formula you employ in (a) to obtain electron speed for an collector potential of 10 MV. Do you see what is wrong? In what way is the formula to be modified?

Solution 20:

- (a) Potential difference across the evacuated tube, $V = 500 \text{ V}$ Specific charge of an electron, e/m
 $= 1.76 \times 10^{11} \text{ C kg}^{-1}$

The speed of each emitted electron is given by the relation for kinetic energy as:

Therefore, the speed of each emitted electron is

$$KE = \frac{1}{2}mv^2 = eV$$

$$\begin{aligned} \therefore v &= \left(\frac{2eV}{m} \right)^{\frac{1}{2}} = \left(2V \times \frac{e}{m} \right)^{\frac{1}{2}} \\ &= \left(2 \times 500 \times 1.76 \times 10^{11} \right)^{\frac{1}{2}} = 1.366 \times 10^7 \text{ m/s} \end{aligned}$$

- (b) Potential of the anode, $V = 10 \text{ MV} = 10 \times 10^6 \text{ V} = 10^7 \text{ V}$ The speed of each electron is given as:

$$\begin{aligned}
 v &= \left(2V \frac{e}{m} \right)^{\frac{1}{2}} \\
 &= \left(2 \times 10^7 \times 1.76 \times 10^{11} \right)^{\frac{1}{2}} \\
 &= 1.88 \times 10^9 \text{ m/s}
 \end{aligned}$$

This result is wrong because nothing can move faster than light. In the above formula, the expression $(mv^2/2)$ for energy can only be used in the non-relativistic limit, i.e., for $v \ll c$.

For very high speed problems, relativistic equations must be considered for solving them. In the relativistic limit, the total energy is given as:

$$E = mc^2 = m_0 \left(1 - \frac{v^2}{c^2} \right)^{-\frac{1}{2}} \text{ Where,}$$

m = Relativistic mass

m_0 = Mass of the particle at rest Kinetic energy is given as:

$$K = mc^2 - m_0c^2$$

Question 21:

(a) A mono energetic electron beam with electron speed of $5.20 \times 10^6 \text{ m s}^{-1}$ is subject to a magnetic field of $1.30 \times 10^{-4} \text{ T}$ normal to the beam velocity. What is the radius of the circle traced by the beam, given e/m for electron equals $= 1.76 \times 10^{11} \text{ Ckg}^{-1}$.

(b) Is the formula you employ in (a) valid for calculating radius of the path of a 20 MeV electron beam? If not, in what way is it modified?

[Note: Exercises 11.20(b) and 11.21(b) take you to relativistic mechanics which is beyond the scope of this book. They have been inserted here simply to emphasis the point that the formulas you use in part (a) of the exercises are not valid at very high speeds or energies. See answers at the end to know what ‘very high speed or energy’ means.]

Solution 21:

(a) Speed of an electron, $v = 5.20 \times 10^6$ m/s

Magnetic field experienced by the electron, $B = 1.30 \times 10^{-4}$ T Specific charge of an electron, $e/m = 1.76 \times 10^{11}$ C kg⁻¹ Where,

$e =$ Charge on the electron = 1.6×10^{-19} C $m =$ Mass of the electron = 9.1×10^{-31} kg⁻¹

The force exerted on the electron is given as:

$$F = e|\vec{v} \times \vec{B}|$$

$$= evB \sin \theta$$

$\theta =$ Angle between the magnetic field and the beam velocity. The magnetic field is normal to the direction of beam.

$$\theta = 90^\circ$$

$$F = evB \quad \dots\dots\dots(1)$$

The beam traces a circular path of radius, r . It is the magnetic field, due to its bending

nature, that provides the centripetal force $\left(F = \frac{mv^2}{r} \right)$ for the beam. Hence, equation (1) reduces

to:

$$E = \frac{1}{2}mv^2$$

$$v = \left(\frac{2E}{m} \right)^{\frac{1}{2}}$$

$$= \sqrt{\frac{2 \times 20 \times 10^6 \times 1.6 \times 10^{-19}}{9.1 \times 10^{-31}}} = 2.652 \times 10^9 \text{ m/s}$$

$$evB = \frac{mv^2}{r}$$

$$r = \frac{mv}{eB} = \frac{v}{\left(\frac{e}{m}\right)B}$$

$$= \frac{5.20 \times 10^6}{(1.76 \times 10^{11}) \times 1.30 \times 10^{-4}} = 0.227 \text{ m} = 22.7 \text{ cm}$$

Therefore, the radius of the circular path is 22.7 cm.

(b) Energy of the electron beam, $E = 20 \text{ M eV} = 20 \times 10^6 \times 1.6 \times 10^{-19} \text{ J}$ The energy of the electron is given as:

This result is incorrect because nothing can move faster than light. In the above formula, the expression $(mv^2/2)$ for energy can only be used in the non-relativistic limit, i.e., for $v \ll c$.

When very high speeds are concerned, the relativistic domain comes into consideration. In the relativistic domain, mass is given as:

$$m = m_0 \left[1 - \frac{v^2}{c^2} \right]^{-\frac{1}{2}}$$

Where,

m_0 = Mass of the particle at rest

Hence, the radius of the circular path is given as: $r = mv / eB$

$$= \frac{m_0 v}{eB \sqrt{\frac{c^2 - v^2}{c^2}}}$$

Question 22:

An electron gun with its collector at a potential of 100 V fires out electrons in a spherical bulb containing hydrogen gas at low pressure (10^{-2} mm of Hg). A magnetic field of $2.83 \times 10^{-4} \text{ T}$ curves the path of the electrons in a circular orbit of radius 12.0 cm. (The path can be viewed because the gas ions in the path focus the beam by attracting electrons, and emitting light by electron capture; this method is known as the ‘fine beam tube’ method. Determine e/m from the data.

Solution 22:

Potential of an anode, $V = 100 \text{ V}$

Magnetic field experienced by the electrons, $B = 2.83 \times 10^{-4} \text{ T}$ Radius of the circular orbit $r =$

$$12.0 \text{ cm} = 12.0 \times 10^{-2} \text{ m}$$

Mass of each electron = m , Charge on each electron = e , Velocity of each electron = v

The energy of each electron is equal to its kinetic energy, i.e.,

$$\frac{1}{2}mv^2 = eV$$

$$v^2 = \frac{2eV}{m} \quad \dots\dots\dots(1)$$

It is the magnetic field, due to its bending nature, that provides the centripetal force

$\left(F = \frac{mv^2}{r} \right)$ for the beam. Hence, we can write:

Centripetal force = Magnetic force

$$\frac{mv^2}{r} = evB$$

$$eB = \frac{mv}{r}$$

$$v = \frac{eBr}{m}$$

Putting the value of v in equation (1), we get:

$$\frac{2eV}{m} = \frac{e^2 B^2 r^2}{m^2}$$

$$\frac{e}{m} = \frac{2V}{B^2 r^2}$$

$$= \frac{2 \times 100}{(2.83 \times 10^{-4})^2 \times (12 \times 10^{-2})^2} = 1.73 \times 10^{11} \text{ C kg}^{-1}$$

Therefore, the specific charge ratio (e/m) is

$$1.73 \times 10^{11} \text{ C kg}^{-1}$$

Question 23:

(a) An X-ray tube produces a continuous spectrum of radiation with its short wavelength end at 0.45 \AA . What is the maximum energy of a photon in the radiation?

Chapter 11

Dual Nature of Radiation and Matter

(b) From your answer to (a), guess what order of accelerating voltage (for electrons) is required in such a tube?

Solution 23:

(a) Wavelength produced by an X-ray tube, $\lambda = 0.45 \text{ \AA} = 0.45 \times 10^{-10} \text{ m}$, Planck's constant,
 $h = 6.626 \times 10^{-34} \text{ Js}$

Speed of light, $c = 3 \times 10^8 \text{ m/s}$

The maximum energy of a photon is given as:

$$E = \frac{hc}{\lambda}$$

$$= \frac{6.625 \times 10^{-34} \times 3 \times 10^8}{0.45 \times 10^{-10} \times 1.6 \times 10^{-19}}$$

$$= 27.6 \times 10^3 \text{ eV} = 27.6 \text{ keV}$$

Therefore, the maximum energy of an X-ray photon is 27.6 k eV.

(b) Accelerating voltage provides energy to the electrons for producing X-rays. To get an X-ray of 27.6 keV, the incident electrons must possess at least 27.6 keV of kinetic electric energy. Hence, an accelerating voltage of the order of 30 keV is required for producing X-rays.

Question 24:

In an accelerator experiment on high-energy collisions of electrons with positrons, a certain event is interpreted as annihilation of an electron-positron pair of total energy

10.2 BeV into two γ -rays of equal energy. What is the wavelength associated with each γ -ray?

(1BeV = 10^9 eV)

Solution 24:

Total energy of two γ -rays: $E = 10.2 \text{ BeV}$

$$= 10.2 \times 10^9 \text{ eV}$$

$$= 10.2 \times 10^9 \times 1.6 \times 10^{-19} \text{ J}$$

Hence, the energy of each γ -ray:

$$E' = \frac{E}{2}$$

$$= \frac{10.2 \times 1.6 \times 10^{-10}}{2} = 8.16 \times 10^{-10} \text{ J}$$

Plank's constant, $h = 6.626 \times 10^{-34} \text{ Js}$

Speed of light $c = 3 \times 10^8 \text{ m/s}$

Energy is related to wavelength as:

$$E' = \frac{hc}{\lambda}$$

$$\lambda = \frac{hc}{E'}$$

$$= \frac{6.626 \times 10^{-34} \times 3 \times 10^8}{8.16 \times 10^{-10}} = 2.436 \times 10^{-16} \text{ m}$$

Therefore, the wavelength associated with each γ -ray is $2.436 \times 10^{-16} \text{ m}$

Question 25:

Estimating the following two numbers should be interesting. The first number will tell you why radio engineers do not need to worry much about photons! The second number tells you why our eye can never 'count photons', even in barely detectable light.

- (a) The number of photons emitted per second by a Medium wave transmitter of 10 kW power, emitting radio waves of wavelength 500 m.
- (b) The number of photons entering the pupil of our eye per second corresponding to the minimum intensity of white light that we humans can perceive ($\sim 10^{-10} \text{ W m}^{-2}$). Take the area of the pupil to be about 0.4 cm^2 , and the average frequency of white light to be about $6 \times 10^{14} \text{ Hz}$.

Solution 25:

(a) Power of the medium wave transmitter, $P = 10 \text{ kW} = 10^4 \text{ W} = 10^4 \text{ J/s}$ Hence, energy emitted by the transmitter per second, $E = 10^4$, Wavelength of the radio wave, $\lambda = 500 \text{ m}$

The energy of the wave is given as:

$$E_1 = \frac{hc}{\lambda}$$

Where, $h = \text{Planck's constant} = 6.6 \times 10^{-34} \text{ Js}$ $c = \text{Speed of light} = 3 \times 10^8 \text{ m/s}$

$$E_1 = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{500} = 3.96 \times 10^{-28} \text{ J}$$

Let n be the number of photons emitted by the transmitter.

$$\therefore n E_1 = E$$

$$n = \frac{E}{E_1}$$

$$= \frac{10^4}{3.96 \times 10^{-28}} = 2.525 \times 10^{31}$$

$$= 3 \times 10^{34}$$

The energy (E_1) of a radio photon is very less, but the number of photons (n) emitted per second in a radio wave is very large.

The existence of a minimum quantum of energy can be ignored and the total energy of a radio wave can be treated as being continuous.

(b) Intensity of light perceived by the human eye, $I = 10^{-10} \text{ W m}^{-2}$ Area of a pupil,

$$A = 0.4 \text{ cm}^2 = 0.4 \times 10^{-4} \text{ m}^2$$

Frequency of white light, $\nu = 6 \times 10^{14} \text{ Hz}$. The energy emitted by a photon is given as: $E = h\nu$

Where,

$$h = \text{Planck's constant} = 6.6 \times 10^{-34} \text{ Js}, E = 6.6 \times 10^{-34} \times 6 \times 10^{14}$$

$$= 3.96 \times 10^{-19} \text{ J}$$

Let n be the total number of photons falling per second, per unit area of the pupil. The total energy per unit for n falling photons is given as:

$$E = n \times 3.96 \times 10^{-19} \text{ J s}^{-1} \text{ m}^{-2}$$

The energy per unit area per second is the intensity of light.

$$E = I$$

$$n \times 3.96 \times 10^{-19} = 10^{-10}$$

$$n = \frac{10^{-10}}{3.96 \times 10^{-19}}$$

$$= 2.52 \times 10^8 \text{ m}^2 \text{ s}^{-1}$$

The total number of photons entering the pupil per second is given as: $n_A = n \times A$

$$= 2.52 \times 10^8 \times 0.4 \times 10^{-4}$$

$$= 1.008 \times 10^4 \text{ s}^{-1}$$

This number is not as large as the one found in problem (a), but it is large enough for the human eye to never see the individual photons.

Question 26:

Ultraviolet light of wavelength 2271 \AA from a 100 W mercury source irradiates a photocell made of molybdenum metal. If the stopping potential is -1.3 V , estimate the work function of the metal. How would the photo-cell respond to a high intensity (10^5 W m^{-2}) red light of wavelength 6328 \AA produced by a He-Ne laser?

Solution 26:

Wavelength of ultraviolet light, $\lambda = 2271 \text{ \AA} = 2271 \times 10^{-10} \text{ m}$ Stopping potential of the metal, $V_0 = 1.3 \text{ V}$

Planck's constant, $h = 6.6 \times 10^{-34} \text{ J}$

Charge on an electron, $e = 1.6 \times 10^{-19} \text{ C}$

Work function of the metal, $= \phi_0$ Frequency of light $= \nu$

We have the photo-energy relation from the photoelectric effect as: $\phi_0 = h\nu - eV_0$

$$= \frac{hc}{\lambda} - eV_0$$

Let ν_0 be the threshold frequency of the metal.

$$\phi_0 = h\nu_0$$

$$\nu_0 = \frac{\phi_0}{h}$$

$$= \frac{6.64 \times 10^{-19}}{6.6 \times 10^{-34}} = 1.006 \times 10^{15} \text{ Hz}$$

Wavelength of red light $\lambda_c = 6328 \text{ \AA}$

Frequency of red light, $\nu_r = \frac{c}{\lambda_c} = 6328 \times 10^{-10} \text{ m}$

$$= \frac{3 \times 10^8}{6328 \times 10^{-10}} = 4.74 \times 10^{14} \text{ Hz}$$

Since $\nu_0 > \nu_r$, the photocell will not respond to the red light produced by the laser.

Question 27:

Monochromatic radiation of wavelength 640.2 nm ($1\text{nm} = 10^{-9}\text{ m}$) from a neon lamp irradiates photosensitive material made of caesium on tungsten. The stopping voltage is measured to be 0.54 V. The source is replaced by an iron source and its 427.2 nm line irradiates the same photo-cell. Predict the new stopping voltage.

Solution 27:

Wavelength of the monochromatic radiation, $\lambda = 640.2\text{ nm} = 640.2 \times 10^{-9}\text{ m}$

Stopping potential of the neon lamp, $V_0 = 0.54\text{ V}$ Charge on an electron, $e = 1.6 \times 10^{-19}\text{ C}$

Planck's constant, $h = 6.6 \times 10^{-34}\text{ Js}$

Let ϕ_0 be the work function and ν be the frequency of emitted light. We have the photo-energy relation from the photoelectric effect as:

$$\begin{aligned}
 eV_0 &= h\nu - \phi_0 \\
 \phi_0 &= \frac{hc}{\lambda} - eV_0 \\
 &= \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{640.2 \times 10^{-9}} - 1.6 \times 10^{-19} \times 0.54 \\
 &= 3.093 \times 10^{-19} - 0.864 \times 10^{-19} \\
 &= 2.229 \times 10^{-19}\text{ J} \\
 &= \frac{2.229 \times 10^{-19}}{1.6 \times 10^{-19}} = 1.39\text{ eV}
 \end{aligned}$$

Wavelength of the radiation emitted from an iron source, $\lambda' = 427.2\text{ nm}$

$$= 427.2 \times 10^{-9}\text{ m}$$

Let V_0' be the new stopping potential. Hence, photo-energy is given as:

$$\begin{aligned}
 eV_0' &= \frac{hc}{\lambda'} = \phi_0 \\
 &= \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{427.2 \times 10^{-9}} - 2.229 \times 10^{-19} \\
 &= 4.63 \times 10^{-19} - 2.229 \times 10^{-19} \\
 &= 2.401 \times 10^{-19} \text{ J} \\
 &= \frac{2.401 \times 10^{-19}}{1.6 \times 10^{-19}} = 1.5 \text{ eV}
 \end{aligned}$$

Hence, the new stopping potential is 1.50 eV.

Question 28:

A mercury lamp is a convenient source for studying frequency dependence of photoelectric emission, since it gives a number of spectral lines ranging from the UV to the red end of the visible spectrum. In our experiment with rubidium photo-cell, the following lines from a mercury source were used: $\lambda_1 = 3650 \text{ \AA}$, $\lambda_2 = 4047 \text{ \AA}$, $\lambda_3 = 4358 \text{ \AA}$, $\lambda_4 = 5461 \text{ \AA}$, $\lambda_5 = 6907 \text{ \AA}$. The stopping voltages, respectively, were measured to be: $V_{01} = 1.28 \text{ V}$, $V_{02} = 0.95 \text{ V}$, $V_{03} = 0.74 \text{ V}$, $V_{04} = 0.16 \text{ V}$, $V_{05} = 0 \text{ V}$.

Determine the value of Planck's constant h , the threshold frequency and work function for the material.

[Note: You will notice that to get h from the data, you will need to know e (which you can take to be $1.6 \times 10^{-19} \text{ C}$). Experiments of this kind on Na, Li, K, etc. were performed by Millikan, who, using his own value of e (from the oil-drop experiment) confirmed Einstein's photoelectric equation and at the same time gave an independent estimate of the value of h .]

Solution 28:

Einstein's photoelectric equation is given as:

$$\begin{aligned}
 eV_0 &= h\nu - \phi_0 \\
 V_0 &= \frac{h}{e}\nu - \frac{\phi_0}{e} \quad \dots\dots\dots(1)
 \end{aligned}$$

Where,

V_0 = Stopping potential, h = Planck's constant

e = Charge on an electron, ν = Frequency of radiation

Chapter 11

Dual Nature of Radiation and Matter

ϕ_0 = Work function of a material

It can be concluded from equation (1) that potential V_0 is directly proportional to frequency ν .

$$\nu = \frac{\text{Speed of light}(c)}{\text{Wavelength}(\lambda)}$$

Frequency is also given by the relation: This relation can be used to obtain the frequencies of the various lines of the given wavelengths.

$$\nu_1 = \frac{c}{\lambda_1} = \frac{3 \times 10^8}{3650 \times 10^{-10}} = 8.219 \times 10^{14} \text{ Hz}$$

$$\nu_2 = \frac{c}{\lambda_2} = \frac{3 \times 10^8}{4047 \times 10^{-10}} = 7.412 \times 10^{14} \text{ Hz}$$

$$\nu_3 = \frac{c}{\lambda_3} = \frac{3 \times 10^8}{4358 \times 10^{-10}} = 6.884 \times 10^{14} \text{ Hz}$$


$$\nu_4 = \frac{c}{\lambda_4} = \frac{3 \times 10^8}{5461 \times 10^{-10}} = 5.493 \times 10^{14} \text{ Hz}$$

$$\nu_5 = \frac{c}{\lambda_5} = \frac{3 \times 10^8}{6907 \times 10^{-10}} = 4.343 \times 10^{14} \text{ Hz}$$

The given quantities can be listed in tabular form as:

Frequency $\times 10^{14}$ Hz	8.219	7.412	6.884	5.493	4.343
Stopping potential V_0	1.28	0.95	0.74	0.16	0

The following figure shows a graph between ν and V_0 .



It can be observed that the obtained curve is a straight line. It intersects the ν -axis at 5×10^{14} Hz, which is the threshold frequency (ν_0) of the material. Point D corresponds to a frequency less than the threshold frequency. Hence, there is no photoelectric emission for the λ_5 line, and therefore, no stopping voltage is required to stop the current.

$$\text{Slope of the straight line} = \frac{AB}{CB} = \frac{1.28 - 0.16}{(8.124 - 5.493) \times 10^{14}}$$

From equation (1), the slope $\frac{h}{e}$ can be written as:

$$\frac{h}{e} = \frac{1.28 - 0.16}{(8.124 - 5.493) \times 10^{14}}$$

$$h = \frac{1.12 \times 1.6 \times 10^{-19}}{2.726 \times 10^{14}} \\ = 6.573 \times 10^{-34} \text{ Js}$$

The work function of the metal is given as:

$$\phi_0 = hv_0 \\ = 6.573 \times 10^{-34} \times 5 \times 10^{14} \\ = 3.286 \times 10^{-19} \text{ J} \\ = \frac{3.286 \times 10^{-19}}{1.6 \times 10^{-18}} \\ = 2.054 \text{ eV}$$

Question 29:

The work function for the following metals is given:

Na = 2.75 eV; K = 2.30 eV; Mo = 4.17 eV; Ni = 5.15 eV. Which of these metals will not give photoelectric emission for a radiation of wavelength 3300 Å from a He-Cd laser placed 1 m away from the photocell? What happens if the laser is brought nearer and placed 50 cm away?

Solution 29:

Mo and Ni will not show photoelectric emission in both cases Wavelength for a radiation,

$$\lambda = 3300 \text{ \AA} = 3300 \times 10^{-10} \text{ m}, \text{ Speed of light, } c = 3 \times 10^8 \text{ m/s}$$

$$\text{Planck's constant, } h = 6.6 \times 10^{-34} \text{ Js}$$

The energy of incident radiation is given as:

$$\begin{aligned}
 E &= \frac{hc}{\lambda} \\
 &= \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{3300 \times 10^{-10}} = 6 \times 10^{-19} \text{ J} \\
 &= \frac{6 \times 10^{-19}}{1.6 \times 10^{-19}} = 3.158 \text{ eV}
 \end{aligned}$$

It can be observed that the energy of the incident radiation is greater than the work function of Na and K only. It is less for Mo and Ni. Hence, Mo and Ni will not show photoelectric emission. If the source of light is brought near the photocells and placed 50 cm away from them, then the intensity of radiation will increase. This does not affect the energy of the radiation. Hence, the result will be the same as before. However, the photoelectrons emitted from Na and K will increase in proportion to intensity.

Question 30:

Light of intensity 10^{-5} W m^{-2} falls on a sodium photo-cell of surface area 2 cm^2 . Assuming that the top 5 layers of sodium absorb the incident energy, estimate time required for photoelectric emission in the wave-picture of radiation. The work function for the metal is given to be about 2 eV. What is the implication of your answer?

Solution 30:

Intensity of incident light, $I = 10^{-5} \text{ W m}^{-2}$

Surface area of a sodium photocell, $A = 2 \text{ cm}^2 = 2 \times 10^{-4} \text{ m}^2$ Incident power of the light,

$$\begin{aligned}
 P &= I \times A \\
 &= 10^{-5} \times 2 \times 10^{-4} \\
 &= 2 \times 10^{-9} \text{ W}
 \end{aligned}$$

Work function of the metal, $\phi_0 = 2 \text{ eV}$

$$\begin{aligned}
 &= 2 \times 1.6 \times 10^{-19} \\
 &= 3.2 \times 10^{-19} \text{ J}
 \end{aligned}$$

Number of layers of sodium that absorbs the incident energy, $n = 5$

Chapter 11

Dual Nature of Radiation and Matter

We know that the effective atomic area of a sodium atom, A_e is 10^{-20} m^2 .

Hence, the number of conduction electrons in n layers is given as:

$$\begin{aligned} n' &= n \times \frac{A}{A_e} \\ &= 5 \times \frac{2 \times 10^{-4}}{10^{-20}} = 10^{17} \end{aligned}$$

The incident power is uniformly absorbed by all the electrons continuously. Hence, the amount of energy absorbed per second per electron is:

$$\begin{aligned} E &= \frac{P}{n'} \\ &= \frac{2 \times 10^{-9}}{10^{17}} = 2 \times 10^{-26} \text{ J/s} \end{aligned}$$

Time required for photoelectric emission:

$$\begin{aligned} t &= \frac{\phi_0}{E} \\ &= \frac{3.2 \times 10^{-19}}{2 \times 10^{-26}} = 1.6 \times 10^7 \text{ s} = 0.507 \text{ years} \end{aligned}$$

The time required for the photoelectric emission is nearly half a year, which is not practical.

Hence, the wave picture is in disagreement with the given experiment.

Question 31:

Crystal diffraction experiments can be performed using X-rays, or electrons accelerated through appropriate voltage. Which probe has greater energy? (For quantitative comparison, take the wavelength of the probe equal to 1 \AA , which is of the order of inter-atomic spacing in the lattice)

$$(m_e = 9.11 \times 10^{-31} \text{ kg}) .$$

Solution 31:

An X-ray probe has a greater energy than an electron probe for the same wavelength.

Wavelength of light emitted from the probe, $\lambda = 1 \text{ \AA} = 10^{-10} \text{ m}$

Mass of an electron, $m_e = 9.11 \times 10^{-31} \text{ kg}$ Planck's constant, $h = 6.6 \times 10^{-34} \text{ Js}$ Charge on an

electron, $e = 1.6 \times 10^{-19} \text{ C}$

The kinetic energy of the electron is given as:

$$E = \frac{1}{2} m_e v^2$$

$$m_e v = \sqrt{2Em_e}$$

Where,

v = Velocity of the electron

$m_e v$ = Momentum (p) of the electron

According to the de Broglie principle, the de Broglie wavelength is given as:

$$\lambda = \frac{h}{m_0 v} = \frac{h}{\sqrt{2Km_0}}$$

$$E = \frac{h^2}{2\lambda^2 m_e}$$

$$= \frac{(6.6 \times 10^{-34})^2}{2 \times (10^{-10})^2 \times 9.11 \times 10^{-31}} = 2.39 \times 10^{-17}$$

$$= \frac{2.39 \times 10^{-17}}{1.6 \times 10^{-19}} = 149.375 eV$$

Energy of photon $E' = \frac{hc}{\lambda a} eV$

$$= \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{10^{-10} \times 1.6 \times 10^{-19}}$$

$$= 12.375 \times 10^3 eV = 12.375 keV$$

Hence, a photon has a greater energy than an electron for the same wavelength.

Question 32:

(a) Obtain the de Broglie wavelength of a neutron of kinetic energy 150 eV. As you have seen in Exercise 11.31, an electron beam of this energy is suitable for crystal diffraction experiments. Would a neutron beam of the same energy be equally suitable? Explain.

$$(m_n = 1.675 \times 10^{-27} kg)$$

(b) Obtain the de Broglie wavelength associated with thermal neutrons at room temperature (27 °C). Hence explain why a fast neutron beam needs to be thermalised with the environment before

it can be used for neutron diffraction experiments.

Solution 32:

(a) De Broglie wavelength 2.327×10^{-12} m; neutron is not suitable for the diffraction experiment

Kinetic energy of the neutron, $K = 150$ eV

$$= 150 \times 1.6 \times 10^{-19}$$

$$= 2.4 \times 10^{-17} \text{ J}$$

Mass of a neutron, $m_n = 1.675 \times 10^{-27} \text{ kg}$

The kinetic energy of the neutron is given by the relation:

$$K = \frac{1}{2} m_0 v^2$$

$$m_0 v = \sqrt{2Km_0}$$

Where,

v = Velocity of the neutron $m_0 v$ = Momentum of the neutron

De-Broglie wavelength of the neutron is given as:

$$\lambda = \frac{h}{m_0 v} = \frac{h}{\sqrt{2Km_0}}$$

$$\lambda = \frac{6.6 \times 10^{-34}}{\sqrt{2 \times 2.4 \times 10^{-17} \times 1.675 \times 10^{-27}}}$$

$$= 2.327 \times 10^{-2} \text{ m}$$

It is clear that wavelength is inversely proportional to the square roots of mass.

Hence, wavelength decrease with increase in mass and vice-versa.

It is given in the previous problem that the inter-atomic spacing of a crystal is about 1 \AA , i.e., 10^{-10} m. Hence, the inter-atomic spacing is about a hundred times greater. Hence, a neutron beam of energy

150 eV is not suitable for diffraction experiments.

(b) De Broglie wavelength = 1.447×10^{-16} m; Room temperature, $T = 27^\circ\text{C} = 27 + 273 = 300$ K

The average kinetic energy of the neutron is given as:

$$E = \frac{3}{2} kT$$

Where,

$$k = \text{Boltzmann constant} = 1.38 \times 10^{-23} \text{ J mol}^{-1} \text{ K}^{-1} .$$

The wavelength of the neutron is given as:

$$\lambda = \frac{h}{m_0 v} = \frac{h}{\sqrt{2Km_0}}$$

$$\begin{aligned} \lambda &= \frac{6.6 \times 10^{-34}}{\sqrt{3 \times 1.675 \times 10^{-27} \times 1.38 \times 10^{-23} \times 300}} \\ &= 1.447 \times 10^{-10} \text{ m} \end{aligned}$$

This wavelength is comparable to the inter-atomic spacing of a crystal. Hence, the high energy neutron beam should first be thermalised, before using it for diffraction.

Question 33:

An electron microscope uses electrons accelerated by a voltage of 50 kV. Determine the de Broglie wavelength associated with the electrons. If other factors (such as numerical aperture, etc.) are taken to be roughly the same, how does the resolving power of an electron microscope compare with that of an optical microscope which uses yellow light?

Solution 33:

Electrons are accelerated by a voltage, $V = 50 \text{ kV} = 50 \times 10^3 \text{ V}$ Charge on an electron, $e = 1.6 \times 10^{-19} \text{ C}$

Mass of an electron, $m_e = 9.11 \times 10^{-31} \text{ kg}$ Wavelength of yellow light = $5.9 \times 10^{-7} \text{ m}$. The

kinetic energy of the electron is given as: $E = eV$

$$= 1.6 \times 10^{-19} \times 50 \times 10^3$$

$$= 8 \times 10^{-15} \text{ J}$$

De Broglie wavelength is given by the relation:

$$\begin{aligned}\lambda &= \frac{h}{\sqrt{2m_c E}} \\ &= \frac{6.6 \times 10^{-34}}{\sqrt{2 \times 9.11 \times 10^{-31} \times 8 \times 10^{-15}}} \\ &= 5.467 \times 10^{-12} \text{ m}\end{aligned}$$

This wavelength is nearly 10^5 times less than the wavelength of yellow light.

The resolving power of a microscope is inversely proportional to the wavelength of light used.

Thus, the resolving power of an electron microscope is nearly 10^5 times that of an optical microscope.

Question 34:

The wavelength of a probe is roughly a measure of the size of a structure that it can probe in some detail. The quark structure of protons and neutrons appears at the minute length- scale of 10^{-15} m or less. This structure was first probed in early 1970's using high energy electron beams produced by a linear accelerator at Stanford, USA. Guess what might have been the order of energy of these electron beams. (Rest mass energy of electron = 0.511 M eV)

Solution 34:

Wavelength of a proton or a neutron, $\lambda \approx 10^{-15}$ m Rest mass energy of an electron:

$$\begin{aligned}m_0 c^2 &= 0.511 \text{ M eV} \\ &= 0.511 \times 10^6 \times 1.6 \times 10^{-19} \\ &= 0.8176 \times 10^{-13} \text{ J}\end{aligned}$$

Planck's constant, $h = 6.6 \times 10^{-34}$ Js Speed of light, $c = 3 \times 10^8$ m/s

The momentum of a proton or a neutron is given as

$$\begin{aligned}p &= \frac{h}{\lambda} \\ &= \frac{6.6 \times 10^{-34}}{10^{-15}} = 6.6 \times 10^{-19} \text{ kg m/s}\end{aligned}$$

The relativistic relation for energy (E) is given as:

$$\begin{aligned}
 E^2 &= p^2 c^2 + m_0^2 c^4 \\
 &= (6.6 \times 10^{-19} \times 3 \times 10^8)^2 + (0.8176 \times 10^{-23})^2 \\
 &= 392.04 \times 10^{-22} + 0.6685 \times 10^{-26} \\
 &= 392.04 \times 10^{-22}
 \end{aligned}$$

$$\begin{aligned}
 E &= 1.98 \times 10^{-10} \text{ J} \\
 &= \frac{1.98 \times 10^{-10}}{1.6 \times 10^{-19}} \\
 &= 1.24 \times 10^9 \text{ eV} = 1.24 \text{ BeV}
 \end{aligned}$$

Thus, the electron energy emitted from the accelerator at Stanford, USA might be of the order of 1.24 B eV.

Question 35:

Find the typical de Broglie wavelength associated with a He atom in helium gas at room temperature (27 °C) and 1 atm pressure; and compare it with the mean separation between two atoms under these conditions.

Solution 35:

De Broglie wavelength associated with He atom = $0.7268 \times 10^{-10} \text{ m}$ Room temperature, $T = 27^\circ\text{C}$
 $= 27 + 273 = 300 \text{ K}$

Atmospheric pressure, $P = 1 \text{ atm} = 1.01 \times 10^5 \text{ Pa}$ Atomic weight of a He atom = 4

Avogadro's number, $N_A = 6.023 \times 10^{23}$ Boltzmann constant, $k = 1.38 \times 10^{-23} \text{ J mol}^{-1} \text{ K}^{-1}$

Average energy of a gas at temperature T, is given as:

$$E = \frac{3}{2} kT$$

De Broglie wavelength is given by the relation:

$$\lambda = \frac{h}{\sqrt{2mE}}$$

Where,

$m =$ Mass of a He atom

$$= \frac{\text{Atomic weight}}{N_A}$$

$$\begin{aligned}
 &= \frac{4}{6.023 \times 10^{23}} \\
 &= 6.64 \times 10^{-24} \text{ g} = 6.64 \times 10^{-27} \text{ kg} \\
 \lambda &= \frac{h}{\sqrt{3mkT}} \\
 &= \frac{6.6 \times 10^{-34}}{\sqrt{3 \times 6.64 \times 10^{-27} \times 1.38 \times 10^{-23} \times 300}} \\
 &= 0.7268 \times 10^{-10} \text{ m}
 \end{aligned}$$

We have the ideal gas formula: $PV = RT$

$$PV = kNT$$

$$\frac{V}{N} = \frac{kT}{P}$$

Where,

V = Volume of the gas

N = Number of moles of the gas

Mean separation between two atoms of the gas is given by the relation:

$$\begin{aligned}
 r &= \left(\frac{V}{N} \right)^{\frac{1}{3}} = \left(\frac{kT}{P} \right)^{\frac{1}{3}} \\
 &= \left[\frac{1.38 \times 10^{-23} \times 300}{1.01 \times 10^5} \right]^{\frac{1}{3}} \\
 &= 3.35 \times 10^{-9} \text{ m}
 \end{aligned}$$

Hence, the mean separation between the atoms is much greater than the de Broglie wavelength.

Question 36:

Compute the typical de Broglie wavelength of an electron in a metal at 27 °C and compare it with the mean separation between two electrons in a metal which is given to be about 2×10^{-10} m.

[Note: Exercises 11.35 and 11.36 reveal that while the wave-packets associated with gaseous molecules under ordinary conditions are non-overlapping, the electron wave packets in a metal strongly overlap with one another. This suggests that whereas molecules in an ordinary gas can be distinguished apart, electrons in a metal cannot be distinguished apart from one another. This indistinguishability has many fundamental implications which you will explore in more advanced Physics courses.]

Solution 36:

Temperature, $T = 27^\circ\text{C} = 27 + 273 = 300\text{ K}$

Mean separation between two electrons, $r = 2 \times 10^{-10}\text{ m}$ De Broglie wavelength of an electron is given as:

$$\lambda = \frac{h}{\sqrt{3mKT}}$$

Where,

$h = \text{Planck's constant} = 6.6 \times 10^{-34}\text{ Js}$

$m = \text{Mass of an electron} = 9.11 \times 10^{-31}\text{ kg}$

$k = \text{Boltzmann constant} = 1.38 \times 10^{-23}\text{ J mol}^{-1}\text{ K}^{-1}$

$$\begin{aligned}\lambda &= \frac{6.6 \times 10^{-34}}{\sqrt{3 \times 9.11 \times 10^{-31} \times 10^{23} \times 300}} \\ &= 6.2 \times 10^{-9}\text{ m}\end{aligned}$$

Hence, the de Broglie wavelength is much greater than the given inter-electron separation.

Question 37:

Answer the following questions:

(a) Quarks inside protons and neutrons are thought to carry fractional charges $[(+2/3)e ; (-1/3)e]$.

Why do they not show up in Millikan's oil-drop experiment?

(b) What is so special about the combination e/m ? Why do we not simply talk of e and m separately?

(c) Why should gases be insulators at ordinary pressures and start conducting at very low pressures?

(d) Every metal has a definite work function. Why do all photoelectrons not come out with the same energy if incident radiation is monochromatic? Why is there an energy distribution of photoelectrons?

(e) The energy and momentum of an electron are related to the frequency and wavelength of the associated matter wave by the relations:

$$E = h\nu, p = \frac{h}{\lambda}$$

Chapter 11

Dual Nature of Radiation and Matter

But while the value of λ is physically significant, the value of v (and therefore, the value of the phase speed $v\lambda$) has no physical significance. Why?

Solution 37:

(b) Quarks inside protons and neutrons carry fractional charges. This is because nuclear force increases extremely if they are pulled apart. Therefore, fractional charges may exist in nature; observable charges are still the integral multiple of an electrical charge.

(b) The basic relations for electric field and magnetic field are

$$\left(eV = \frac{1}{2}mv^2 \right) \text{ and } \left(eBv = \frac{mv^2}{r} \right) \text{ respectively}$$

These relations include e (electric charge), v (velocity), m (mass), V (potential), r (radius), and B (magnetic field). These relations give the value of velocity of an electron as

$$\left(v = \sqrt{2V \left(\frac{e}{m} \right)} \right) \text{ and } \left(v = Br \left(\frac{e}{m} \right) \right)$$

It can be observed from these relations that the dynamics of an electron is determined not by e and m separately, but by the ratio e/m .

(c) At atmospheric pressure, the ions of gases have no chance of reaching their respective electrons because of collision and recombination with other gas molecules. Hence, gases are insulators at atmospheric pressure. At low pressures, ions have a chance of reaching their respective electrodes and constitute a current. Hence, they conduct electricity at these pressures.

(d) The work function of a metal is the minimum energy required for a conduction electron to get out of the metal surface. All the electrons in an atom do not have the same energy level. When a ray having some photon energy is incident on a metal surface, the electrons come out from different levels with different energies. Hence, these emitted electrons show different energy distributions.

(e) The absolute value of energy of a particle is arbitrary within the additive constant. Hence, wavelength (λ) is significant, but the frequency (ν) associated with an electron has no direct physical significance.

Therefore, the product $\nu \lambda$ (phase speed) has no physical significance. Group speed is given as:

$$\begin{aligned}v_G &= \frac{dv}{dk} \\ &= \frac{dv}{d\left(\frac{1}{\lambda}\right)} = \frac{dE}{dP} = \frac{d\left(\frac{p^2}{2m}\right)}{dP} = \frac{p}{m}\end{aligned}$$

This quantity has a physical meaning.
